

## LITERATURE CITED

- 41<sup>st</sup> Northeast Regional Stock Assessment Workshop. 2006 (revised and reissued August 2006). 41<sup>st</sup> SAW assessment summary report. U.S. Dept. Commerce Northeast Fisheries Science Center Ref. Doc 05-10:36p.
- 48<sup>th</sup> Northeast Regional Stock Assessment Workshop. 2009. 48<sup>th</sup> SAW assessment summary report. U.S. Dept. Commerce Northeast Fisheries Science Center Ref. Doc 09-10:395p.
- Acosta, A, and R.S. Appeldoorn. 1992. Estimation of growth, mortality and yield per recruit for *Lutjanus synagris* (Linnaeus) in Puerto Rico. Bull. Mar. Sci. 50:282–291.
- Abitia-Cardenas, L.A., F. Galvan-Magaña, and J. Rodriguez-Romero. 1997. Food habits and energy values of prey of striped marlin, *Tetrapterus audax*, off the east coast of Mexico. Fish. Bull. 95:360–368.
- Ahrenholz, D.W. 1991. Population biology and life history of the North American menhadens, *Brevoortia* spp. Mar. Fish. Rev. 53(4):3–19.
- Allen, D. M. 1979. Biological aspects of the calico scallop, *Argopecten gibbus*, determined from spat monitoring. The Nautilus 93:107–119.
- Andovora, F., and C. Pipitone. 1997. Food and feeding habits of the amberjack, *Seriola dumerili*, in the central Mediterranean Sea during the spawning season. Cahiers de Biologie Marine 38:9196.
- Armitage, T.M., and W.S. Alevizon. 1980. The diet of Florida pompano (*Trachinotus carolinus*) along the east coast of central Florida. Fla. Sci. 43:19–26.
- Armstrong, M.P., and R.G. Muller. 1996. A summary of biological information for southern kingfish (*Menticirrhus americanus*), gulf kingfish (*M. littoralis*), and northern kingfish (*M. saxatilis*) in Florida waters. Report from the Florida Marine Research Institute to the Florida Marine Fisheries Commission. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1996-004)
- Armstrong, M.P., R.E. Crabtree, M.D. Murphy, and R.G. Muller. 1996a. A stock assessment of tripletail, *Lobotes surinamensis*, in Florida waters. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1996-001)
- Armstrong, M.P., P.B. Hood, M.D. Murphy, and R.G. Muller. 1996b. A stock assessment of permit, *Trachinotus falcatus*, in Florida waters. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1996-005)

Armstrong, M.P., M.D. Murphy, R.G. Muller, D. Harshany, and R.E. Crabtree. 1996c. A stock assessment of hardhead catfish, *Arius felis*, and gafftopsail catfish, *Bagre marinus*, in Florida waters. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee, Florida. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR 1996-003)

Arnold, W.S. 1986. Review summary of the hard clam, *Mercenaria* spp., in Florida. Florida Department of Natural Resources, Bureau of Marine Research, St. Petersburg, FL., unpublished manuscript.

Arnold, W.S. 1995. Summary report on the calico scallop (*Argopecten gibbus*) fishery of the southeastern United States. Unpublished manuscript, Florida Department of Environmental Protection, Florida Marine Research Institute, St. Petersburg, FL.

Arnold, W.S, T.M. Bert, D.C. Marelli, H. Cruz-Lopez, and P.A. Gill. 1996. Genotype-specific growth of hard clams (genus *Mercenaria*) in a hybrid zone: variation among habitats. *Marine Biology* 125:129–139.

Arnold, W.S., S.P. Geiger, S.P. Stephenson. 2009. *Mercenaria mercenaria* introductions into Florida, USA, waters: duration, not size of introduction, influences genetic outcomes. *Aquat. Biol.* 5: 49-62.

Arnold, W.S., D.C. Marelli, T.M. Bert, D.S. Jones, and I.R. Quitmyer. 1991. Habitat-specific growth of hard clams *Mercenaria mercenaria* (L.) from the Indian River, Florida. *J. Exp. Mar. Biol. Ecol.* 147:245–265.

Arnold, W.S., S.L. Walters, S.J. Fajans, S.C. Peters, T.M. Bert. 2004. Influence of congeneric aquaculture on hard clam (*Mercenaria* spp.) population genetic structure. *Aquac. Int.* 12: 139-160

Arnoldi, D.C., W.H. Herke, and E.J. Clairain, Jr. 1974. Estimates of growth rate and length of stay in a marsh nursery of juvenile Atlantic croaker, *Micropogonias undulatus* (Linnaeus), 'sandblasted' with fluorescent pigments. *Proc. Gulf and Carib. Fish. Inst.* 26:158–172.

Arrenguin-Sanchez, F., M.A Cabrera, and F.A. Aguilar. 1995. Population dynamics of the king mackerel (*Scomberomorus cavalla*) of the Campeche Bank, Mexico. *Sci. Mar.* 59:637–645.

Atlantic States Marine Fisheries Commission. 1982. Fishery management plan for the summer flounder (*Paralichthys dentatus*) fishery. Fisheries Management Report No. 3. Washington, D.C.

Atlantic States Marine Fisheries Commission. 2005. Atlantic croaker stock assessment and peer review reports: 2003 Atlantic croaker stock assessment report, 2003 terms of reference and advisory report, 2004 Atlantic croaker stock assessment report, 2004 terms of reference and advisory report. Atlantic States Marine Fisheries Commission, Washington, D.C. 370p.

Atlantic States Marine Fisheries Commission. 2010. Spot Life History Report. Report to the ASMFC South Atlantic State/Federal Fisheries Management Board. April 2010. 66p. Available at: <http://www.asmf.org/uploads/file/spotLifeHistoryReportApr10.pdf>

Atlantic States Marine Fisheries Commission. 2014. Black Drum stock assessment. Stock Assessment Report No. 14-02. 331 p.

Atlantic States Marine Fisheries Commission. 2015. Bluefish benchmark stock assessment for 2015. Atlantic States Marine Fisheries Commission, Stock assessment report, 536 p.

Atlantic States Marine Fisheries Commission. 2016. Weakfish stock assessment and peer review report. Atlantic States Marine Fisheries Commission, Stock assessment report, 435 p.

Atlantic States Marine Fisheries Commission. 2017. Spot stock assessment and peer review report. Atlantic States Marine Fisheries Commission, Stock assessment report, 435 p.

Austin, H.M., D. Scoles, and A.J. Abell. 1999. Morphometric separation of annual cohorts within mid-Atlantic bluefish, *Pomatomus saltatrix*, using discriminant function analysis. Fish. Bull. 97:411-420.

Babcock, L.L. 1951. The tarpon, 5<sup>th</sup> edition. Privately Printed, Buffalo, N.Y. 157p.

Bagley, M.J., D.G. Lindquist, and J.B. Geller. 1999. Microsatellite variation, effective population size, and population genetic structure of vermilion snapper, *Rhomboplites aurorubens*, off the southeastern USA. Marine Biology 134:609–620.

Barber, R.C. 1989. Age and growth of vermilion snapper (*Rhomboplites aurorubens*) in the northern Gulf of Mexico. M.S. Thesis. Texas A&M University. 87p.

Barbieri, L.Z., M.E. Chittenden, Jr., and C.M. Jones. 1994. Age, growth, and mortality of Atlantic croaker, *Micropogonias undulatus*, in the Chesapeake Bay region, with a discussion of apparent geographic changes in population dynamics. Fish. Bull., U.S. 92:1–12.

Barbieri, L.Z., M.E. Chittenden, Jr., and C.M. Jones. 1997. Yield-per-recruit analysis and management strategies for Atlantic croaker, *Micropogonias undulatus*, in the Middle Atlantic Bight. Fish. Bull., U.S. 95:637–645.

Barger, L.E. 1985. Age and growth of Atlantic croakers in the northern Gulf of Mexico, based on otolith sections. Trans. Amer. Fish. Soc. 114:847–850.

Barger, L.E. 1990. Age and growth of bluefish *Pomatomus saltatrix* from the northern Gulf of Mexico and U.S. south Atlantic coast. Fish. Bull., U.S.:88(4):805–809.

Bearden, C.M. 1963. A contribution to the biology of the king whittings, genus *Menticirrhus*, of South Carolina. Contributions from Bear Bluff Laboratories 38.

Beardsely, G.L., Jr. 1967. Age, growth, and reproduction of the dolphin, *Coryphaena hippurus*, in the Straits of Florida. Copeia 1967:441–451.

Beardsley, G.L. and W. Richards. 1970. Size, seasonal abundance and length-weight relation of some scombrid fishes from southeast Florida. U.S. Fish. Wildl. Serv. Spec. Sci. Rep. 595:1–5.

- Beaumariage, D.S. 1969. Current status of biological investigations of Florida's mackerel fisheries. Proc. Gulf Caribb. Fish. Inst. 22nd Annual Meeting, 1969:79–86.
- Beaumariage, D.S. 1973. Age, growth and reproduction of king mackerel, *Scomberomorus cavalla*, in Florida. Fla. Mar. Res. Publ. 1. 45p.
- Beckman, D.W., A.L. Stanley, J.H. Render, C.A. Wilson. 1990. Age and growth of black drum in Louisiana waters of the Gulf of Mexico. Trans. Amer. Fish. Soc. 119:537–544.
- Beckman, D.W., A.L. Stanley, J.H. Render, C.A. Wilson. 1991. Age and growth-rate estimation of sheepshead *Archosargus probatocephalus* in Louisiana waters using otoliths. Fish. Bull., U.S. 89:1–8.
- Behringer, D.C., M.J. Butler IV, J.D. Shields, J. Moss. 2011. Review of *Panulirus argus* virus 1—a decade after its discovery. Diseases of Aquatic Organisms 94:153-160. doi: 10.3354/dao02326
- Bellinger, J.W., and J.W. Avault, Jr. 1971. Food habits of juvenile pompano, *Trachinotus carolinus*, in Louisiana. Trans. Amer. Fish. Soc. 100:486–494.
- Benson, N.G. (ed.). 1982. Life history requirements of selected finfish and shellfish in Mississippi Sound and adjacent waters. U.S. Fish Wildl. Serv., Biol. Rep. FWS/OBS-81/51, 97p.
- Bentivoglio, A.A. 1988. Investigations into the growth, maturity, mortality rates, and occurrence of the dolphin (*Coryphaena hippurus* Linnaeus) in the Gulf of Mexico. M.S. Thesis, University College of North Wales, Bangor, U.K., 37p.
- Ben-Tuvia, A. 1960. Synopsis of biological data on *Sardinella aurita* of the Mediterranean Sea and other waters. FAO Fish. Biol. Synop. 14:287–312.
- Berkeley, S.A., and E. D. Houde. 1978. Biology of two exploited species of halfbeaks, *Hemiramphus brasiliensis* and *H. balao* from southeast Florida. Bull. Mar. Sci. 28:624–644.
- Berkeley, S.A., and E.D. Houde. 1984. Age, growth, and mortality of coastal herrings in the eastern Gulf of Mexico. Final Report to the Gulf and South Atlantic Development Foundation, Inc. Proj. 21-12-32272.
- Berkeley, S.A., E.D. Houde, and F. Williams. 1975. Fishery and biology of ballyhoo on the southeast Florida coast. Univ. Miami Sea Grant Special Report 4.
- Berrigan, M. 2000. Oyster resource assessment report. Apalachicola Bay 1999–2000. Department of Agriculture and Consumer Services. Division of Aquaculture. Unpaged.
- Berrigan, M., T. Candies, J. Cirino, R. Dugas, C. Dyer, J. Gray, T. Herrington, W. Keithly, R. Leard, J.R. Nelson, M. Van Hoose. 1991. The oyster fishery of the Gulf of Mexico, United States: a regional management plan. Gulf States Marine Fisheries Commission Publ. No. 24.

- Berry, F.H. 1959. Young jack crevalles (*Caranx* species) off the southeastern Atlantic coast of the United States. *Fish. Bull.* 59:417–535.
- Berry, F.H., and I. Barrett. 1963. Gillraker analysis and speciation in the thread herring genus *Opisthonema*. *Inter-Am. Trop. Tuna Comm. Bull.* 7:111–190.
- Berry, F.H., and E.S. Iverson. 1967. Pompano: biology, fisheries, and farming potential. *Proc. 19th Annual Session Gulf and Caribb. Fish. Inst.* 116–128.
- Berry, F.H., and W.F. Smith-Vanez. 1978. Carangidae, *FAO Species Identification Sheets for Fisheries Purposes. Western Central Atlantic*. Ed. W. Fisher, FAO, Rome. Volume II.
- Bert, T.M. 1992. Summary of workshop on current issues related to stone crab (genus *Menippe*) biology and fisheries, p. 108–115. *In*: T.M. Bert (editor), *Proceedings of a symposium on stone crab (genus Menippe) biology and fisheries*. Florida Mar. Res. Pub. 50.
- Bert, T.M., and R.G. Harrison. 1988. Hybridization in western Atlantic stone crabs (genus *Menippe*): evolutionary history and ecological context influence species interactions. *Evolution* 42:528–544.
- Bert, T.M., and D.M. Hesselman, W.S. Arnold, W.S. Moor, H. Cruz-Lopez, and D.C. Marelli. 1993. High frequency of gonadal neoplasia in a hard clam (*Mercenaria* spp.) hybrid zone. *Marine Biology* 117: 97-104.
- Bertelsen, R.D. and J.H. Hunt. 1992. Results of the 1991 mail surveys of recreational lobster fishermen (Special Sport Season and Regular Season Surveys). Florida Department of Environmental Protection. Florida Marine Research Institute. Unpublished manuscript.
- Bigelow, H.B., and W.C. Schroeder. 1953. *Fishes of the Gulf of Maine*. U.S. Fish Wildl. Serv. Fish. Bull. 53.
- Bohnsack, J.A. and D.E. Harper. 1988. Length-weight relationships of selected marine reef fishes from the southeastern United States and the Caribbean. *NOAA Tech. Mem.NMFS-SEFC-215*:31p.
- Boothby, R.N., and W.J. Avault, Jr. 1971. Food habits, length-weight relationship, and condition factor of the red drum (*Sciaenops ocellata*) in southeastern Louisiana. *Trans. Amer. Fish. Soc.* 100(2):290–295.
- Bortone, S.A., and J.L. Williams. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (South Florida)—gray, lane, mutton, and yellowtail snappers. U.S. Fish and Wildl. Serv. Biol. Rep. 82(11.52). U.S. Army Corps of Engineers, TR EL-82-4. 18p.
- Bortone, S.A., R.N. Besser, C.R. McNeil. 1997. The importance of Silas Stearns (1859-1888) and the Pensacola red snapper fishery to North American Ichthyology. *Collection Building in Ichthyology and Herpetology*. Chap. 25, pp 459-467. American Society of Ichthyologists and Herpetologists.
- Bradley, E., and C.E. Bryan. 1975. Life history and fishery of the red snapper (*Lutjanus campechanus*) in the northwestern Gulf of Mexico: 1970–1974. *Proc. Gulf Caribb. Fish. Inst.* 17:77–106.

- Breuer, J.P. 1957. An ecological survey of Baffin and Alazan bays, Texas. *Publ. Inst. Mar. Sci.* 4:134–155.
- Brule, T., C., T. Deniel, t. Colas-Marrufo, and M. Sanchez-Crespo. 1999. Red grouper reproduction in the southern Gulf of Mexico. *Trans. Amer. Fish. Soc.* 128:385–402.
- Buckel, J.A., M.J. Fogarty, and D.O. Conover. 1999a. Foraging habits of bluefish, *Pomatomus saltatrix*, on the U.S. east coast continental shelf. *Fish. Bull.* 97:758–775.
- Buckel, J.A., M.J. Fogarty, and D.O. Conover. 1999b. Mutual prey of fish and humans: a comparison of biomass consumed by bluefish, *Pomatomus saltatrix*, with that harvested by fisheries. *Fish. Bull.* 97:776–785.
- Bullock, L.H., and G.B. Smith. 1991. Seabasses (Pisces: Serranidae). *Memoirs of the Hourglass Cruises Vol. VIII, Part II.*
- Burch, R.K. 1979. The greater amberjack, *Seriola dumerili*: Its biology and fishery off southeastern Florida. MS Thesis, University of Miami, Coral Gables, FL.
- Burns, K.M. 1981. Seasonal and areal distribution of scombrid larvae in the vicinity of Palm Beach, Florida. M.A. Thesis, University of South Florida, Tampa. 66p.
- Burns, K.M., C. Neidig, J. Lotz, and R. Overstreet. 1998. Cobia (*Rachycentron canadum*, stock assessment study in the Gulf of Mexico and in the South Atlantic. Final report for NOAA-NMFS MARFIN Award No. NA57FF0294, St. Petersburg, Florida.
- Buroker, N.E. 1983. Population genetics of the American oyster, *Crassostrea virginica*, along the Atlantic coast and the Gulf of Mexico. *Marine Biology* 75:99–112.
- Burton, M.L. 2001. Age, growth and mortality of gray snapper, *Lutjanus griseus*, from the east coast of Florida. *Fish. Bull.* 99:254–265.
- Butler, C.B. and T.R. Matthews. 2015. Effects of ghost fishing lobster traps in the Florida Keys. *ICES Journal of Marine Science*. doi: 10.1093/icesjms/fsu238.
- Caldwell, D.K. 1965. Systematics and variation in the sparid fish *Archosargus probatocephalus*. *Bull. South Calif. Acad. Sci.* 64:89–100.
- Camber, C.I. 1955. A survey of the red snapper fishery of the Gulf of Mexico, with special reference to the Campeche Banks. *Fla. Board Conserv. Mar. Res. Lab. Tech. Ser.* 12. 64p.
- Campana, S.E., and C.M. Jones. 1998. Radiocarbon from nuclear testing applied to age validation of black drum, *Pogonias cromis*. *Fish. Bull.* 96:185–192.

- Camper, J.D., R.C. Barber, L.R. Richardson, and J.R. Gold. 1993. Mitochondrial DNA variation among red snapper (*Lutjanus campechanus*) from the Gulf of Mexico. *Molecular Marine Biology and Biotechnology* 2:154–161.
- Campton, D.E., and B. Mahmoudi. 1991. Allozyme variation and population structure of striped mullet (*Mugil cephalus*) in Florida. *Copeia* 1991:485–492.
- Carles, C. 1967. Algunos datos sobre la biología del banano, *Elops saurus* L. (Teleostomi: Elopidae). *Centr. Invest. Pesq., Inst. Nacl. Pesca, Cuba, Contrib.* 27:1–53.
- Carr, W.E.S. and C.A. Adams. 1972. Food habits of juvenile marine fishes: evidence of the cleaning habit in the leatherjacket, *Oligoplites saurus*, and the spottail pinfish, *Diplodus holbrooki*. *Fish. Bull., U.S.* 70(4):1111–1120.
- Chagaris, D., D. Addis, and B. Mahmoudi. 2014. The 2014 stock assessment update for striped mullet, *Mugil cephalus*, in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute. 76 pp.
- Chagaris, D., B. Mahmoudi, and M. D. Murphy. 2015. The 2015 stock assessment of Red Drum, *Sciaenops ocellatus*, in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2015-003: 95 p.
- Chagaris, D., B. Mahmoudi, M. D. Murphy, and C. Guenther. 2012. Status of flounder fishery resources in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2012-xxx: 74 p.
- Chapman, R.W., G.R. Sedberry, C.C. Koenig, and B.M. Eleby. 1999. Stock identification of gag, *Mycteroperca microlepis*, along the southeast coast of the United States. *Mar. Biotechnol.* 1:137–146.
- Chikhi, L., F. Bonhomme, and J. F. Agnese. 1998. Low genetic variability in a widely distributed and abundant clupeid species, *Sardinella aurita*. New empirical results and interpretations. *J. Fish Biol.* 52:861–878.
- Christmas, J. Y., Jr. and G. Gunter 1960. Distribution of menhaden, genus *Brevoortia*, in the Gulf of Mexico. *Trans. Amer. Fish. Soci.* 89(4):338–343.
- Claro, R., A. Garcia-Cagide and R. Fernandez de Albiza. 1989. Características biológicas del pez perro, *Lachnolaimus maximus* (Walbaum) en el Golfo de Batabano, Cuba. *Rev. Invest. Mar.* 10(3):239–252.
- Clifton, K.B. and P.J. Motta. 1998. Feeding morphology, diet, and ecomorphological relationships among five Caribbean labrids (Teleostei, Labridae). *Copeia* 1998:953–966.
- Cody, R.P., J.R. O’Hop, S. Brown, and L.A. Hallock. 2000. Final report on at-sea observations of pompano netting (August 1999 – June 2000). Report to the Florida Fish and Wildlife Conservation Commission from the Florida Marine Research Institute, St. Petersburg.

- Coleman, F.C., C.C. Koenig, W.F. Herrnkind. 1993. Survey of Florida inshore shrimp by-catch. Second Annual Report, Florida Department of Natural Resources, Florida Marine Research Institute, Contract C-7779. 30p.
- Coleman, F.C., C.C. Koenig, and L.A. Collins. 1996. Reproductive styles of shallow-water grouper (Pisces: Serranidae) in the eastern Gulf of Mexico and the consequences of fishing spawning aggregations. *Environ. Biol. Fishes* 47:129–141.
- Colin, P.L. 1982. Spawning and larval development of the hogfish, *Lachnolaimus maximus* (Pisces: Labridae). *Fish. Bull. U.S.* 80:853–862.
- Collete, B.B., and C.E. Nauen. 1983. FAO species catalog. Vol2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. FAO Fish. Synop. 125, 137p.
- Collins, L.A. and J.H. Finucane. 1989. Reproductive biology of yellowtail snapper, *Ocyurus chrysurus*, from the Florida Keys. U.S. Department of Commerce, NOAA, NMFS, SEFSC, Panama City Laboratory, PCL Contrib. No. 89-11, 23 p.
- Collins, L.A., A.G. Johnson, C.C. Koenig, and M.S Baker, Jr. 1998. Reproductive patterns, sex ratio, and fecundity in gag, *Mycteroperca microlepis* (Serranidae), a protogynous grouper from the northeastern Gulf of Mexico. *Fish. Bull., U.S.* 96:415–427.
- Collins, L.A., G.R. Fitzhugh, L.A. Lombardi-Carlson, H.M. Lyon, W.T. Walling, and D.W. Oliver. 2002. Characterization of red grouper (Serranidae: *Epinephelus morio*) reproduction from the eastern Gulf of Mexico. NOAA, NMFS Panama City Laboratory Contribution Series 2002-07. 20p.
- Collins MR, Stender BW. 1987. Larval king mackerel (*Scomberomorus cavalla*), Spanish mackerel (*S. maculatus*), and bluefish (*Pomatomus saltatrix*) off the southeast coast of the United States, 1973-1980. *Bulletin of Marine Science* 41:822-834.
- Collins, M.R., C.W. Waltz, W.A. Roumillat, and D.L. Stubbs. 1987. Contribution to the life history and reproductive biology of gag, *Mycteroperca microlepis* (Serranidae), in the South Atlantic Bight. U.S. Natl. Mar. Fish. Serv. *Fish. Bull.* 85:648–653.
- Cook, H.L., and M.A. Murphy. 1965. Early development stages of the rock shrimp, *Sicyonia brevirostrus* Stimpson, reared in the laboratory. *Tulane Stud. Zoology* 12:109–127.
- Cooper, W., A. Collins, J. O’Hop, and D. Addis. 2014. SEDAR 37 - The 2013 Stock assessment report for hogfish in the South Atlantic and Gulf of Mexico. FWC-FWRI. 295 pp.
- Cooper, W., R. Gandy, and C. Crowley. 2013. A stock assessment for blue crab, *Callinectes sapidus*, in Florida waters through 2011. Report to the Florida Fish and Wildlife Commission Division of Marine Fisheries Management. Florida Fish and Wildlife Research Institute In House Report 2013-xxx. 127 p.

- Cordes, J.F. and J.E. Graves. 2003. Investigation of congeneric hybridization in and stock structure of weakfish (*Cynoscion regalis*) inferred from analysis of nuclear and mitochondrial DNA loci. *Fish. Bull.* 101:443-450.
- Costello, T., T.M. Bert, D.G. Cartano, G. Davis, G. Lyons, C. Rockwood, J. Stevely, J. Tashiro, W.L. Trent, D. Turgeon, and J. Zuboy. 1979. Fishery management plan for the stone crab fishery in the Gulf of Mexico. *Federal Register* 44: 19445–19496.
- Cox, C., J.H. Hunt, W.G. Lyons, and G.E. Davis. 1997. Nocturnal foraging of the Caribbean spiny lobster (*Panulirus argus*) on offshore reefs of Florida, USA. *Mar. Freshwater Res.* 48:671–679.
- Crabtree, R.E. 1995. Relationship between lunar phase and spawning activity of tarpon, *Megalops atlanticus*, with notes on the distribution of larvae. *Bull. Mar. Sci.* 56:895–899.
- Crabtree, R.E., E.C. Cyr, D.C. Chaverri, W.O. McLarney, and J.M. Dean. 1997. Reproduction of tarpon, *Megalops atlanticus*, from Florida and Costa Rican waters and notes on their age and growth. *Bull. Mar. Sci.* 62:271–285.
- Crabtree, R.E., E.C. Cyr, and J.M. Dean. 1995. Age and growth of tarpon, *Megalops atlanticus*, from South Florida waters. *Fish. Bull., U.S.* 93:619–628.
- Crabtree, R.E., C. Harnden, and D. Snodgrass. 1997. Maturation and reproductive seasonality in bonefish, *Albula vulpes*, from the waters of the Florida Keys. *Fish. Bull., U.S.* 456–465.
- Crabtree, R.E., C. Harnden, D. Snodgrass, C. Stevens. 1996. Age and growth of bonefish, *Albula vulpes*, from the waters of the Florida Keys. *Fish. Bull., U.S.* 94:442–451.
- Crabtree, R.E., P.B. Hood, and D. Snodgrass. 2002. Age, growth, and reproduction of permit, *Trachinotus falcatus*, in Florida waters. *Fish. Bull.* 100:26–34.
- Crabtree, R.E., C. Stevens, D. Snodgrass, F.J. Stengard. 1998. Feeding habits of bonefish, *Albula vulpes*, from the waters of the Florida Keys. *Fish. Bull., U.S.*:754–766.
- Crawford, M.K., C.B. Grimes, and N.E. Buroker. 1989. Stock identification of weakfish, *Cynoscion regalis*, in the middle Atlantic region. *Fish. Bull.* 87:205-211.
- Crales, M.M., and T.N. Lee. 1995. Larval distribution and transport of penaeiid shrimps during the presence of the Tortugas Gyre in May–June 1991. *Fish. Bull., U.S.* 93:471–482.
- Crosetti, D., W.S. Nelson, and J.C. Avise. 1994. Pronounced genetic structure of mitochondrial DNA among populations of the circumglobally distributed grey mullet (*Mugil cephalus*). *J. Fish Biology* 44:47–58.
- Cummings, N.J., and D.B. McClellan. 1996. Movement patterns and stock interchange of greater amberjack, *Seriola dumerili* in the southeastern United States. U.S. Dep. Comm. NOAA NMFS SFSC Miami Laboratory Contr. Rep. MIA-95/96-14.

- Cummings, N.J., and D.B. McClellan. 1998. An analysis of the Gulf of Mexico greater amberjack, *Seriola dumerili*, stock condition. Proc. Gulf Carib. Fish. Inst. 50:206–227.
- Dahlberg, M.D. 1970. Atlantic and Gulf of Mexico menhadens, genus *Brevoortia* (Pisces:Clupeidae). Bull. Fla. State Mus. Biol. Sci. 15:91–162.
- Darcy, G.H. 1983a. Synopsis of biological data on the grunts *Haemulon aurolineatum* and *H. plumieri* (Pisces: Haemulidae). FAO Fisheries Synopsis No. 133.
- Darcy, G.H. 1983b. Synopsis of biological data on the pigfish *Orthopristis chrysoptera* (Pisces: Haemulidae). FAO Fisheries Synopsis No. 134.
- Darcy, G.H. 1985. Synopsis of biological data on the pinfish, *Lagodon rhomboides* (Pisces: Sparidae). FAO Fisheries Synopsis No 141.
- Darnell, R.M. 1958. Food habits of fishes and larger vertebrates of Lake Pontchartrain, Louisiana, an estuarine community. Publ. Inst. Mar. Sci. Univ. Texas 5:354–416.
- Davis, J.C. 1976. Biology of the hogfish, *Lachnolaimus maximus* (Walbaum), in the Florida Keys. Master's Thesis, University of Miami, Coral Gables, Florida.
- Davis, R.L. and Marshall, N. 1961. The feeding of the bay scallop, *Aequipecten irradians*. Proc. Natl. Shellfish. Assoc. 52:25–29.
- Dawson, C.E. 1965. Length-weight relationships of some Gulf of Mexico fishes. Trans. Amer. Fish. Soc. 94:279–280.
- Deegan, L. A. 1985. The population ecology and nutrient transport in gulf menhaden in Fourleague Bay, Louisiana. Doctoral Dissertation. Louisiana State University, Baton Rouge. 134pp.
- De Silva, J.A. 2002. The 2002 update of the quota and stock assessment of bluefish, *Pomatomus saltatrix*, on Florida's Atlantic coast. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2002-003)
- De Silva J.A. and R.G. Muller. 2001. An update of the stock assessment and status of Florida east coast weakfish, *Cynoscion regalis*. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2001–002)
- De Silva, J.A. and M.D. Murphy. 2001. A summary of the status of white grunt, *Haemulon plumieri*, from the east coast of Florida. Florida Fish and Wildlife Conservation Commission Florida Marine Research Institute In-House Report IHR2001-001:18p.

- DeVries, D.A. 1982. Age and growth of spot in North Carolina. Northeast Fish Wildl. Conf., April 13–15, 1982, Cherry Hill, NJ. (Abstract).
- DeVries, D.A., and C.B. Grimes. 1997. Spatial and temporal variation in age and growth of king mackerel, *Scomberomorus cavalla*, 1977–1992. Fish. Bull., U.S. 95:694–708.
- Dick, E. J., MacCall, A.D., 2011. Depletion-Based Stock Reduction Analysis: A catch-based method for determining sustainable yields for data-poor fish stocks. Fish. Res. 110, 331-341.
- Diener, R.A., A. Inglis, and G.B. Adams. 1974. Stomach contents from fishes from Clear Lake and tributary waters, a Texas estuarine area. Contrib. Mar. Sci. 18:7–17.
- Dippold, D. A. 2016. Fishery and Population Dynamics of Mississippi's Spotted Seatrout. *Master's Theses*. 174. [http://aquila.usm.edu/masters\\_theses/174](http://aquila.usm.edu/masters_theses/174).
- Ditty, J.G., R.F. Shaw, C.B. Grimes, and J. S. Cope. 1994. Larval development, distribution, and abundance of common dolphin, *Coryphaena hippurus*, and pompano dolphin, *C. equiselis* (family: Coryphaenidae), in the northern Gulf of Mexico. Fish. Bull., U.S. 92:275–291.
- Doermann, J.E., D. Huddleston, D. Lipsey, and S.H. Thompson. 1977. Age and growth of the sea catfish, *Arius felis*, in Mississippi coastal waters. J. Tenn. Acad. Sci. 52.
- Domeier, M.L., C. Koenig, and F. Coleman. 1996. Reproductive biology of the gray snapper (*Lutjanus griseus*), with notes on spawning for other western Atlantic snappers (Lutjanidae), p. 189–201. In: F. Arreguin-Sanchez, J.L. Munro, M.C. Balgos, and D. Pauly (eds.) Biology and culture of tropical groupers and snappers. ICLARM Conf. Proc. 48, 449p.
- Dutka-Gianelli, J. and D.J. Murie. 2001. Age and growth of sheepshead, *Archosargus probatocephalus* (Pisces: Sparidae), from the northwest coast of Florida. Bull. Mar. Sci. 68:69-83.
- Eldred, B., and W.G. Lyons. 1966. Larval ladyfish, *Elops saurus* Linnaeus 1766 (Elopidae), in Florida and adjacent waters. Fla. Bd. Conserv. Mar. Lab. Leaflet. Ser. 4.
- Epperly, S. P. 1981. A population investigation of Atlantic menhaden: a meristic, morphometric, and biochemical approach. M.S. thesis. University of South Florida, Tampa. 107 pp.
- Fable, W.A., Jr., A.G. Johnson, and L.E. Barger. 1987. Age and growth of Spanish mackerel, *Scomberomorus maculatus*, from Florida and the Gulf of Mexico. Fish. Bull., U.S. 85:777–783.
- Fields, H.M. 1962. Pompanos (*Trachinotus* spp.) of the South Atlantic coast of the United States. U.S. Fish Wildl. Serv., Fish. Bull. 62:189–222.
- Figuerola, M., D. Matos-Caraballo y, and W. Torres. 1998. Maturation and reproductive seasonality of four reef fish species in Puerto Rico. Proceedings of the Gulf and Caribbean Fisheries Institute 50:938-968.

- Finucane, J.H. 1969. Ecology of pompano (*Trachinotus carolinus*) and the permit (*T. falcatus*) in Florida. Trans. Amer. Fish. Soc. 98:478–486.
- Finucane, J.H., C.B. Grimes and S.P. Naughton. 1990. Diets of young king and Spanish mackerel off the Southeast United States. Northeast Gulf Science 11(2):145–153.
- Fisheries-Independent Monitoring Program staff. 2008. Fisheries-independent monitoring program 2007 annual data summary report. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR 2008-006.
- Fitzhugh, G.R., B.A. Thompson, and T. Snider III. 1993. Ovarian development, fecundity, and spawning frequency of black drum *Pogonias cromis* in Louisiana. Fish. Bull., U.S. 91:244–253.
- Fitzhugh, G.R., W.L. Trent, and W.A. Fable, Jr. 1999. Age-structure, mesh-size selectivity, and comparative life history parameters of southern and gulf flounder (*Paralichthys lethostigma* and *P. albigutta*) in northwest Florida. Final Report from the NOAA-NMFS SEFSC Panama City Lab to the Florida Department of Environmental Protection Contract S 3700 831499.
- Forcucci, D., M.J. Butler IV, and J.H. Hunt. 1994. Population dynamics of juvenile Caribbean spiny lobster, *Panulirus argus*, in Florida Bay, Florida. Bull. Mar. Sci. 54:805–818.
- Fore, P.L., and T.W. Schmidt. 1973. Biology of juvenile and adult snook, *Centropomus undecimalis*, in the ten thousand islands. Pages SVI-1–SVI-18. In: Ecosystems analysis of the Big Cypress swamp and estuaries. U.S. Environ. Prot. Agency Surv. Anal. Div., Athens, Ga.
- Franks, J.S., and K.E. VanderKooy. 2000. Feeding habits of juvenile lane snapper *Lutjanus synagris* from Mississippi coastal waters, with comments on the diet of gray snapper *Lutjanus griseus*. Gulf Caribb. Res. 12:11–17.
- Franks, J.S., N. M. Garber, and J.R. Warren. 1996. Stomach contents of juvenile cobia, *Rachycentron canadum*, from the northern Gulf of Mexico. Fish. Bull. 94:374–380.
- Franks, J.S., J.T. McBee, and M.T. Allen. 1992. Studies on the seasonal movements and migratory patterns of cobia, *Rachycentron canadum*, in Mississippi marine waters and adjacent marine waters. Unpublished report, Gulf Coast Research Lab., Ocean Springs, MS 39566.
- Franks, J.S., J.R. Warren, and M.V. Buchanan. 1999. Age and growth of cobia, *Rachycentron canadum*, from the northeastern Gulf of Mexico. Fish. Bull. 97:459–471.
- Fry B., P.L. Mumford, and M.B. Roblee. 1999. Stable isotope studies of pink shrimp (*Farfantepenaeus duorarum*, Burkenroad) migrations on the southwestern Florida shelf. Bull. Mar. Sci. 65:419–430.
- Funicelli, N.A., D.A. Meineke, H.E. Bryant, M.R. Dewey, G.M. Ludwig, and L.S. Mengel. 1989. Movements of striped mullet, *Mugil cephalus*, tagged in Everglades National Park, Florida. Bull. Mar. Sci. 44:1710178.

- Fuss, C.M., Jr., and J.A. Kelly, Jr. 1968. Biology of thread herring. *In*: Sykes, J.E. Report of the Bureau of Fisheries Biological Laboratory, St. Petersburg Beach, Florida, fiscal year 1967. U.S. Fish and Wildl. Circ. 290:14–15.
- Fuss, C.M., Jr., J.A. Kelly, Jr., and K.W. Prest, Jr. 1969. Gulf thread herring: aspects of the developing fishery and biological research. *Proc. Gulf Carib. Fish. Inst.* 21:111–125.
- Galloway, B.J., J.G. Cole, R. Meyer, and P. Roscigno. 1999. Delineation of essential habitat for juvenile red snapper in the northwestern Gulf of Mexico. *Trans. Amer. Fish. Soc.* 128:713–726.
- Galtsoff, P.S. 1964. The American oyster, *Crassostrea virginica* (Gmelin). U.S. Fish and Wildlife Service. *Fish. Bull.* 64.
- Gerhart, S.D., and T.M. Bert. 2008. Life-history aspects of stone crabs (genus *Menippe*): size at maturity, growth, and age. *J. Crust. Biol.* 28:252–261.
- Gibbs, R.H., Jr., and B.B. Collette. 1959. On the identification, distribution, and biology of the dolphins, *Coryphaena hippurus* and *C. equiselis*. *Bull. Mar. Sci.* 9:117–152.
- Gilmore, R.G., C.J. Donohue, and D.W. Cooke. 1983. Observations on the distribution and biology of east-central Florida populations of the common snook, *Centropomus undecimalis* (Bloch). *Fla. Sci. Spec. Suppl. Issue 45(4) Part 2.*
- Gilmore, R.G., Jr., C.J. Donohoe, D.W. Cooke, and D.J. Herrema. 1981. Fishes of the Indian River Lagoon and adjacent waters, Florida. Harbor Branch Found. Tech. Rep. 41.
- Gold, J. R. 2000. Genetic analysis to determine mixing proportions by season of Western Atlantic and Gulf of Mexico stocks of king mackerel. MARFIN Project NA57-FF-0295. Final Report.
- Gold, J.R., and L.R. Richardson. 1991. Genetic studies of marine fishes. IV. An analysis of population structure in the red drum (*Sciaenops ocellatus*) using mitochondrial DNA. *Fish. Res.* 12:213–241.
- Gold, J.R., and L.R. Richardson. 1998a. Population structure in greater amberjack, *Seriola dumerili*, from the Gulf of Mexico and the western Atlantic Ocean. *Fish. Bull.* 96:767–778.
- Gold, J.R., and L.R. Richardson. 1998b. Mitochondrial DNA diversification and population structure in fishes from the Gulf of Mexico and western Atlantic. *J. Heredity* 89:404–414.
- Gold, J.R., A.Y. Kristmundsdottir, and L.R. Richardson. 1997. Mitochondrial DNA variation in king mackerel (*Scomberomorus cavalla*) from the western Atlantic Ocean and Gulf of Mexico. *Marine Biology* 129:221–232.
- Gold, J.R., L.R. Richardson, and C. Furman. 1999. Mitochondrial DNA diversity and population structure of spotted seatrout (*Cynoscion nebulosus*) in coastal waters of the southeastern United States. *Gulf Mex. Sci.* 17:40–50.

- Gold, J.R., F. Sun, and L.R. Richardson. 1997. Population structure of red snapper from the Gulf of Mexico as inferred from analysis of mitochondrial DNA. *Trans. Amer. Fish. Soc.* 126:386–396.
- Gold, J.R., E. Saillant, N.D. Ebel, and S. Lem. 2009. Conservation genetics of gray snapper (*Lutjanus griseus*) in U.S. waters of the northern Gulf of Mexico and western Atlantic Ocean. *Copeia* 2009(2): 277-286.
- Goodyear, C.P. 1995a. Mean size at age: an evaluation of sampling strategies using simulated red grouper data. *Trans. Amer. Fish. Soc.* 124:746–755.
- Goodyear, C.P. 1995b. Red snapper in U.S. waters of the Gulf of Mexico. Draft report dated 10/22/95 (final dated 10/30/95) from NOAA-NMFS SEFSC Miami Lab Contribution MIA-95/96-05.
- Goodyear, C.P. 1996a. Status of the red drum stocks of the Gulf of Mexico. NMFS-SEFSC, Miami Lab. MIA 95/96-47.
- Goodyear, C.P. 1996b. Minimum sizes for red grouper: consequences of considering variable size at age. *Trans. Amer. Fish. Soc.* 16:505–511.
- Goodyear, C.P., and M.J. Schirippa. 1991. A biological profile for vermilion snapper with a description of the fishery in the Gulf of Mexico. NOAA-NMFS SEFSC Miami Laboratory Contribution No. MIA-90/91-78.
- Grall, C. 1984. A study of the biology of the Spanish sardine (*Sardinella aurita*). NOAA Tech. Memo. NMFS-SEFC-187.
- Graves, J.E., J.R. McDowell, A.M. Beardsley, and D.R. Scoles. 1992a. Stock structure of the bluefish *Pomatomus saltatrix* along the mid-Atlantic coast. *Fish. Bull. U.S.* 90:703–710.
- Graves, J.E., J.R. McDowell, M.L. Jones. 1992b. A genetic analysis of weakfish *Cynoscion regalis* stock structure along the mid-Atlantic coast. *Fish. Bull. U.S.* 90:469–475.
- Greccay, P.A., and T.E. Targett. 1996. Spatial patterns in condition and feeding of juvenile weakfish in Delaware Bay. *Trans. Amer. Fish. Soc.* 125:803–808.
- Grimes, C.B., J.H. Finucane, L.A. Collins, and D.A. DeVries. 1990. Young king mackerel *Scomberomorus cavalla*, in the Gulf of Mexico, a summary of the distribution and occurrence of larvae and juveniles, and spawning dates for Mexican juveniles. *Bull. Mar. Sci.* 46:640–654.
- Grimes, C.B., A.G. Johnson, W.A. Fable, Jr. 1987. Delineation of king mackerel, (*Scomberomorus cavalla*), stocks along the U.S. east coast and in the Gulf of Mexico (ABSTRACT). U.S. Dept. Commerce, NOAA Tech. Memo. NMFS\_SEFSC-199:186–187.
- Guillory, V, H. Perry, and S. VanderKooy. 2001. The blue crab fishery of the Gulf of Mexico, United States: a regional management plan. Gulf State Marine Fisheries Commission, Inter-jurisdictional Fisheries Management Program Number 96:301p.

Gulf of Mexico Fishery Management Council. 1981. Fishery management plan for the shrimp fishery of the Gulf of Mexico, United States waters (revised November 1981). Gulf of Mexico Fishery Management Council, Tampa, Florida.

Gulf of Mexico Fishery Management Council's Reef Fish Stock Assessment Panel. 2002. September 2002 report of the reef fish stock assessment panel, final draft. September 17-19, 2002 meeting in Miami, Florida.

Gulf of Mexico and South Atlantic Fishery Management Councils. 1985. Final amendment 1 fishery management plan: environmental impact statement for the coastal pelagic resources (mackerels). Gulf of Mexico Fishery Management Council, Tampa, Florida.

Gunter, G. 1945. Studies on marine fishes of Texas. Publ. Inst. Mar. Sci. 1:1–190.

Haddon, M. 2006. Report on SEDAR 09 for the Centre of Independent Experts, University of Miami. 38pp. National Marine Fisheries Service, SEDAR, Charleston, S.C.

Hales, L.S., and M.J. Van Den Avyle. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (South Atlantic) – spot. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.91). U.S. Army Corps of Engineers TR EL-82-4, 24p.

Hansen, D.J. 1970. Food, growth, migration, reproduction, and abundance of pinfish, *Lagodon rhomboides*, and Atlantic croaker, *Micropogon undulatus*, near Pensacola, Florida, 1963–65. Fish. Bull., U.S. 68:135–146.

Harnden, C.W., R.E. Crabtree, and J.M. Shenker. 1999. Onshore transport of elopomorph leptocephali and glass eels (Pisces: Osteichthyes) in the Florida Keys. Gulf of Mexico Science 1:17–26.

Harrigan, P., J.C. Zieman, and S.A. Macko. 1989. The base of nutritional support for the gray snapper (*Lutjanus griseus*): an evaluation based on the combined stomach contents and stable isotope analysis. Bull. Mar. Sci. 44:65–77.

Harrington, R.W., and E.S. Harrington. 1961. Food selection among fishes invading a high subtropical salt marsh: from onset of flooding through the progress of a mosquito brood. Ecology 42:646–666.

Harris, P.J., D.M. Wyanski, D.B. White, P.P. Mikell, and P.B. Eyo. 2007. Age, growth, and reproduction of greater amberjack off the southeastern U.S. Atlantic coast. Trans. Amer. Fish. Soc. 136:1534-1545.

Hart, R.A. 2012a. Stock assessment of brown shrimp (*Farfantepenaeus aztecus*) in the U.S. Gulf of Mexico for 2011. NOAA Fisheries. Galveston, TX. 43 p.

Hart, R.A. 2012b. Stock assessment of pink shrimp (*Farfantepenaeus duorarum*) in the U.S. Gulf of Mexico for 2011. NOAA Fisheries. Galveston, TX. 17 p.

- Hart, R.A. 2012c. Stock assessment of white shrimp (*Farfantepenaeus setiferus*) in the U.S. Gulf of Mexico for 2011. NOAA Fisheries. Galveston, TX. 41 p.
- Heinemann, D. 2002. Preliminary assessment of bluefish, *Pomatomus saltatrix*, in the Gulf of Mexico. NMFS Southeast Fisheries Science Center, Miami, report submitted to the Gulf Mexico Fisheries Management Council.
- Heist, E.J. and J.R. Gold. 2000. DNA microsatellite loci and genetic structure of red snapper in the Gulf of Mexico. *Trans. Amer. Fish. Soc.* 129:469–475.
- Hensley, V.I. and D.A. Hensley. 1995. Fishes eaten by sooty terns and brown noddies in the Dry Tortugas, Florida. *Bull. Mar. Sci.* 56(3):813–821.
- Hettler, W.F., Jr. 1989. Food habits of juveniles of spotted seatrout and gray snapper in western Florida Bay. *Bull. Mar. Sci.* 44:155–162.
- Hightower, J.E., L. Lee, and P. Rand. 2000. Status of east coast Atlantic croaker – DRAFT 8/17/00. North Carolina State University report to the North Carolina Department of Environment and Natural Resources Division of Marine Fisheries, Morehead City.  
[http://www4.ncsu.edu/unity/users/j/jhncsu/public/ATC\\_StockAssessment.html](http://www4.ncsu.edu/unity/users/j/jhncsu/public/ATC_StockAssessment.html).
- Hildebrand, S.F. 1943. Notes on the affinity, anatomy and development of *Elops saurus* Linnaeus. *J. Wash. Acad. Sci.* 33:90–94.
- Hildebrand, S. F. 1948. A review of the American menhaden, genus *Brevoortia*, with a description of a new species. *Smithsonian Miscellaneous Collection* 107(18):39pp.
- Hildebrand, S.F. 1964. Family Clupeidae. *In*: H.B. Bigelow, (ed.), *Fishes of the western North Atlantic*, p. 257–385. Mem. 1 (part3), Sears Foundation Marine Research, Yale University, New Haven, Connecticut.
- Hood, P.B., and A.K. Johnson. 1997. A study of the age structure, growth, maturity schedules and fecundity of gray triggerfish (*Balistes capriscus*), red porgy (*Pagrus pagrus*), and vermilion snapper (*Rhomboplites aurorubens*) from the eastern Gulf of Mexico MARFIN Grant No. NA57FF0289. Tenth Annual MARFIN Conference, December 10–11, Tampa, FL.
- Hood, P.B., and A.K. Johnson. 1999. Age, growth, mortality, and reproduction of vermilion snapper, *Rhomboplites aurorubens*, from the eastern Gulf of Mexico. *Fish. Bull.*, U.S 97:828–841.
- Hood, P.B., and R.A. Schlieder. 1992. Age, growth, and reproduction of gag, *Mycteroperca microlepis* (Pisces: Serranidae), in the eastern Gulf of Mexico. *Bull. Mar. Sci.* 51:337–352.
- Houde, E.D., C. Grall and S.A. Berkeley. 1983. Population parameter estimates for three shoaling pelagic fishes in the eastern Gulf of Mexico. *Int. Council Explor. Sea, Pelagic Fish Comm.*, C.M. 1983/H:43. 11 p.

- Houde, E.D., J.C. Leak, C.E. Dowd, S.A. Berkeley, and W.J. Richards. 1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. Univ. Miami Rep. BLM Contract. AA550-CT7-28.
- Huntsman, G.R., J. Potts, R. Mays, R.L. Dixon, P.W. Willis, M.L. Burton, and B.W. Harvey. 1992. A stock assessment of the snapper-grouper complex in the U.S. South Atlantic based on fish caught in 1990. Report for South Atlantic Fishery Management Council, Charleston, SC.
- Ibanez Aguirre, A.L., M. Gallardo-Cabello, and X. Chiappa Carrara. 1999. Growth analysis of striped mullet, *Mugil cephalus*, and white mullet, *M. curema* (Pisces: Mugilidae), in the Gulf of Mexico. Fish. Bull., U.S. 97:861–872.
- Johnson, A.G. 1983. Age and growth of yellowtail snapper from South Florida. Trans. Amer. Fish. Soc. 112:173–177.
- Johnson, A.G. and C.H. Saloman. 1984. Age, growth, and mortality of gray triggerfish, *Balistes capriscus* from the northeastern Gulf of Mexico. Fish. Bull. 82:485–492.
- Johnson, A.G. and R. N. Vaught 1986. Species profile of Spanish sardine (*Sardinella aurita*). NOAA Tech. Memo NMFS-SEFC-187. 82p.
- Johnson, A.G., L.A. Collins, J. Dahl, and M.S. Baker, Jr. 1995. Age, growth, and mortality of lane snapper in the northern Gulf of Mexico. Proc. Ann Conf. Southeast Assoc. Fish and Wildl. Agencies 49:178–186.
- Johnson, A.G., L.A. Collins, C.P. Keim. 1994. Age-size structure of gray snapper from the southeastern United States: A comparison of two methods of back-calculating size at age from otoliths. Proc. Annu. Conf. Southeast. Assoc. Fish and Wildl. Agencies 48:592–600.
- Johnson, A.G., W.A. Fable, Jr. C.B. Grimes, L. Trent, and J.V. Perez. 1994. Evidence for distinct stocks of king mackerel, *Scomberomorus cavalla*, in the Gulf of Mexico. Fish. Bull. 92:91–101.
- Johnson, A.G., W.A. Fable, Jr., M.L. Williams, L.E. Barger. 1983. Age, growth, and mortality of king mackerel, *Scomberomorus cavalla*, from the southeastern United States. Fish. Bull., U.S. 81:97–106.
- Johnson, A.K., P. Thomas, and R.R. Wilson, Jr. 1998. Seasonal cycles of gonadal development and plasma sex steroid levels in *Epinephelus morio*, a protogynous grouper in the eastern Gulf of Mexico. J. Fish Biol. 52:502–518.
- Johnson, M.W., S.P. Powers, C.L. Hightower, and M. Kenworthy. 2010. Age, growth, mortality, and diet composition of vermilion snapper from the north-central Gulf of Mexico. Trans. Amer. Fish. Soc. 139:1136-1149.
- Jones, C.M., and B. Wells. 1998. Age, growth, and mortality of black drum, *Pogonias cromis*, in the Chesapeake Bay region. Fish. Bull. 96:451–461.

- Judy, M. H. and R. M. Lewis 1983. Distribution of eggs and larvae of Atlantic menhaden, *Bevoortia tyrannus*, along the Atlantic coast of the United States. U.S. Natl. Mar. Fish. Serv. Spec. Sci. Rep. Fish. 774, 23p.
- Kahn, D., J. Uphoff, V. Crecco, D. Vaughan, B. Murphy, J. Brust, R. O'Reilly, and L. Paramore. 2006. Weakfish assessment report for peer review. A Report by the Stock Assessment Subcommittee to the Weakfish Technical Committee of the Atlantic States Marine Fisheries Commission, February 2006.
- Kahn, D., J. Uphoff, B. Murphy, V. Crecco, J. Brust, R. O'Reilly, L. Paramore, D. Vaughan, and J. de Silva. 2005. A stock assessment of weakfish through 2003. A Report by the Stock Assessment Subcommittee to the Weakfish Technical Committee of the Atlantic States Marine Fisheries Commission, February 4, 2005.
- Karlsson, S., E. Saillant, and J. Gold. 2009. Population structure and genetic variation of lane snapper (*Lutjanus synagris*) in the northern Gulf of Mexico. *Marine Biology* 156:1841-1855.
- Kemp, R.J. 1949. Report on stomach analysis from June 1, 1949 through August 31, 1949. Ann. Rept. Tex Game Fish, Oyster Comm. p. 100–127.
- Kendall, A.W., Jr., and L.A. Walford. 1979. Sources and distribution of bluefish, *Pomatomus saltatrix*, larvae and juveniles off the east coast of the United States. *Fish. Bull.*, U.S.:77(1):213–227.
- Kennedy, F.S., J.J. Crane, R.A.Schlieder, and D.G. Barber. 1977. Studies of the rock shrimp, *Sicyonia brevirostris*, a new fishery resource on Florida's Atlantic shelf. Florida Marine Research Publications 27.
- Killam, K.A. 1992. Synthesis of basic life histories of Tampa Bay fishes. Tampa Bay Estuary Program, Tech. Pub. 10-92, 155p.
- Kinsey, S.T., T. Orsoy, T.M. Bert, and B. Mahmoudi. 1994. Population structure of the Spanish sardine *Sardinella aurita*: natural morphological variation in a genetically homogeneous population. *Marine Biology* 118:309–317.
- Klima, E.F. 1971. Distribution of some coastal pelagic fishes in the western Atlantic. *Comm. Fish. Rev.* 33:21–34.
- Kline, L.L. 1993. A summary of Life history, population dynamics, and yield-per-recruit modeling of Atlantic croaker, *Micropogonias undulatus*, in the Chesapeake Bay area. p. 73–84, *In*: L.L. Cline and H. Speir (editors), Proceedings of a workshop on spot (*Leiostomus xanthurus*) and Atlantic croaker (*Micropogonias undulatus*), October 26–27, 1993, Virginia Institute of Marine Science. Atlantic States Marine Fisheries Commission Special Report No. 25.
- Knapp, F.T. 1949. Menhaden utilization in relation to the conservation of food and game fishes of the Texas gulf coast. *Trans. Amer. Fish. Soc.* 79:137–144.

- Koblinsky, G.J., and P.F. Sheridan. 1979. Distribution, abundance, feeding, and long-term fluctuations of spot, *Leiostomus xanthurus*, and croaker, *Micropogonias undulatus*, in Apalachicola Bay, Florida, 1972–1977. *Contrib. Mar. Sci.* 22:149–161.
- Koenig, C.C. (unpublished manuscript). Final report: spawning biology of shallow-water Gulf of Mexico groupers. MARFIN award No. NA90AA-H-MF748. Available from author at Florida State University, Department of Biological Sciences, Tallahassee, FL 32306-2043.
- Koenig, C.C. 1993. Reproductive biology of gray snapper (*Lutjanus griseus*). Final Report to the Florida Department of Natural Resources, Contract C-7223.
- Kough AS, Paris CB, Butler MJ IV. 2013. Larval Connectivity and the International Management of Fisheries. *PLoS ONE* 8(6): e64970. doi:10.1371/journal.pone.0064970
- Krause, M.K., W.S. Arnold, and W.G. Ambrose, Jr. 1994. Morphological and genetic variation among three populations of calico scallops, *Argopecten gibbus*. *J. Shellfish Res.* 13:529–537.
- Lankford, T.E, Jr., T.E. Targett, and P.M. Gaffney. 1999. Mitochondrial DNA analysis of population structure in the Atlantic croaker, *Micropogonias undulatus* (Perciformes: Sciaenidae). *Fish. Bull.* 97:884–890.
- Larkin, M.F. 2011. Assessment of south Florida’s bonefish stock. Doctoral Dissertation. University of Miami, Coral Gables. 194pp.
- Leard, R., B. Mahmoudi, B. Savoie, H. Blanchet, H. Lazauski, K. Spiller, M. Buchanan, C. Dyer, and W. Keithly. 1995. The striped mullet fishery of the Gulf of Mexico, United States: a regional management plan. Gulf States Marine Fisheries Commission Publication No. 33.
- Leard, R., R. Matheson, K. Meador, W. Keithly, C. Luquest, M. Van Hoose, C. Dyer, S. Gordon, J. Robertson, D. Horn, and R. Scheffler. 1993. The black drum fishery of the Gulf of Mexico, United States: a regional management plan. Gulf States Marine Fisheries Commission No. 28.
- Legault, C.M., and S.C. Turner. 1999. Stock assessment analyses of Atlantic greater amberjack. NMFS-SEFSC Sustainable Fisheries Division Contr. No. SFD-98/99-63.
- Lenihan, H.S., and F. Micheli. 2000. Biological effects of shellfish harvesting on oyster reefs: resolving a fishery conflict by ecological experimentation. *Fish. Bull.*, U.S. 98:86–95.
- Levine, S.J. 1980. Gut contents of forty-four Lake Pontchartrain, Louisiana, fish species. In: Stone, J.H. (ed.), *Environmental analysis of Lake Pontchartrain, Louisiana, its surrounding wetlands, and selected land uses*, Vol. II, p.899–1030. Center for Wetland Resources, Louisiana State University, Baton Rouge, Louisiana.
- Lewis, J.B. 1951. The phyllosoma larvae of the spiny lobster, *Panulirus argus*. *Bull. Mar. Sci. Gulf Caribb.* 1:89–103.

- Lewis, R. M. and C. M. Roithmayer 1981. Spawning and sexual maturity of gulf menhaden *Brevoortia patronus*. Fish. Bull. 78(4):947–951.
- Livingston, R.J. 1984. Trophic response of fishes to habitat variability in coastal seagrass systems. Ecology 65:1258–1275.
- Lombardi-Carlson, L.A., G.R. Fitzhugh, B. Fable, M. Ortiz, and C. Gardner. 2006. Age, length, and growth of Gag (*Mycteroperca microlepis*) from the northeastern Gulf of Mexico: 1979-2005. NOAA, NMFS Panama City Laboratory Contribution 06-03. 57p.
- Lombardi-Carlson, L.A., G.R. Fitzhugh, C. Palmer, C. Gardner, R. Farsky, and M. Ortiz. 2008. Regional size, age and growth differences of red grouper (*Epinephelus morio*) along the west coast of Florida. Fish. Res. 91:239-251.
- Loosanoff, V.L. 1937. Development of the primary gonad and sexual phases of *Venus mercenaria* Linnaeus. Biol. Bull. 72:389–405.
- Lotz, J.M., R. M. Overstreet, and J.S. Franks. 1996. Gonadal maturation in the cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. Gulf Res. Reports 9:147–159.
- Low, R.A., Jr. 1973. Shoreline grassbed fishes in Biscayne Bay, Florida, with notes on availability of clupeid fishes. M.S. Thesis, University Miami, Coral Gables, Florida. 51p.
- Lowerre-Barbieri, S.K., M.E. Chittenden, Jr., and L.R. Barbieri. 1995. Age and growth of weakfish, *Cynoscion regalis*, in the Chesapeake Bay region with a discussion of historical changes in maximum size. Fish. Bull., U.S. 93:643–656.
- Lowerre-Barbieri, S.K., M.E. Chittenden, Jr., and C.M. Jones. 1994. A comparison of a validated otolith method to age weakfish, *Cynoscion regalis*, with the traditional scale method. Fish. Bull., U.S. 92:555–568.
- Lowerre-Barbieri, S.K., S. Villegas-Rios, D. Walters, S. Bickford, J., Cooper, W., Muller, R., Trotter, A. 2014. Spawning site selection and contingent behavior in Common Snook, *Centropomus undecimalis*. PLoS ONE 9(7): e101809. doi:10.1371/journal.pone.0101809
- Luckhurst, B.E., J.M. Dean, and M. Reichert. 2000. Age, growth and reproduction of the lane snapper *Lutjanus synagris* (Pisces: Lutjanidae) at Bermuda. Mar. Ecol. Prog. Ser. 203:255-261.
- Luczkovich, J.J., S.F. Norton, and R.G. Gilmore. 1995. The influence of oral anatomy on prey selection during the ontogeny of two percoid fishes, *Lagodon rhomboides* and *Centropomus undecimalis*. Environ. Biol. Fishes 44:79–95.
- Lukens, R.R. (ed.) 1989. Spanish mackerel fishery management plan. Gulf States Marine Fisheries Commission No. 19, Ocean Springs, Mississippi.
- Lyons, W.G. 1986. Problems and perspectives regarding recruitment of spiny lobster, *Panulirus argus*, to the south Florida fishery. Can. J. Fish. Aquat. Sci. 43:2099–2106.

- MacCall, A.D., 2009. Depletion-corrected average catch: a simple formula for estimating sustainable yields in data-poor situations. *ICES J. Mar. Sci.* 66, 2267-2271.
- Mackerel Stock Assessment Panel. 2003. 2003 report of the mackerel stock assessment panel. Prepared at the Panel Meeting held May 19-21, 2003 in Miami. Gulf of Mexico Fishery Management Council, Tampa, Florida, and South Atlantic Fishery Management Council, Charleston, South Carolina. 31p.
- Mahmoudi, B. 1991. Population assessment of black mullet (*Mugil cephalus*) in the eastern Gulf of Mexico. Final Report of Cooperative Agreement (MARFIN) NA86-WC-H-06138, Florida Department of Environmental Protection, St. Petersburg. 78p.
- Mahmoudi, B. 2000. Status and trends in the Florida mullet fishery and an updated stock assessment. Florida Marine Research Institute report to the Florida Fish and Wildlife Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2000-006)
- Mahmoudi, B. 2005. The 2005 update of the stock assessment for striped mullet, *Mugil cephalus*, in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2005-XXX:43p.
- Mahmoudi, B. 2008. The 2008 update of the stock assessment for striped mullet, *Mugil cephalus*, in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2008-XXX:114p.
- Mahmoudi, B., and R. S. McBride. 2002. A review of Florida's halfbeak bait fishery and halfbeak biology, and a preliminary stock assessment. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2002-015:44p.
- Mahmoudi, B., D. Pierce, and J. O'Hop. 1999. Trends in the Florida baitfish fishery and an update on Spanish sardine stock status along the central west coast. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee, August 1999.
- Mahmoudi, B., D. Pierce, M. Wessel, and R. Lehnert. 2002. Trends in the Florida baitfish fishery and an update on baitfish stock distribution and abundance along the central West Coast of Florida. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee, September 2002. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2002-014)
- Mahmoudi, B., M.D. Murphy, D. Addis. 2010. A stock assessment for pompano, *Trachinotus carolinus*, in Florida waters through 2009. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee.

- Mahon, R. and S. Mahon. 1987. Seasonality and migration of pelagic fishes in the eastern Caribbean. *In* Mahon, R. (ed.), Report and proceedings of the expert consultation on shared fishery resources of the Lesser Antilles region, pp. 192–273. FAO Fisheries Report 383, FAO, Rome.
- Manooch, C.S. III 1976. Age, growth and mortality of the white grunt, *Haemulon plumieri* Lacepède (Pisces: Pomadasyidae), from North Carolina and South Carolina. Proc. Annu. Conf. Southeast. Assoc. Game Fish Comm. 30: 58–70.
- Manooch, C.S. III. 1984. Fisherman's Guide to the Fishes of the Southeastern United States. North Carolina Museum of Natural History, Raleigh, NC.
- Manooch, C.S., III and C.L. Drennon. 1987. Age and growth of yellowtail snapper and queen triggerfish collected from the U.S. Virgin Islands and Puerto Rico. Fish. Res. 6:53-68.
- Manooch, C.S. III, and M. Haimovici. 1978. Age and growth of the gag, *Mycteroperca microlepis*, and size-age composition of the recreational catch off the southeastern United States. Trans. Amer. Fish. Soc. 107:234–240.
- Manooch, C.S. III, and D.L. Mason. 1984. Age, growth, and mortality of lane snapper from southern Florida. Northeast Gulf Sci. 7:109–115.
- Manooch, C.S., III, and R.H. Matheson III. 1981. Age, growth, and mortality of gray snapper collected from Florida waters. Proc. Ann. Conf. S.E. Assoc. Fish Wildl. Agencies 35:331–344.
- Manooch, C.S. III and Potts, J.C. 1997a. Age, growth, and mortality estimates of greater amberjack from the southeastern United States. Fish. Res. 30:229–240.
- Manooch, C.S. III and Potts, J.C. 1997b. Age, growth, and mortality estimates of greater amberjack, *Seriola dumerili*, from the U.S. Gulf of Mexico headboat fishery. Bull. Mar. Sci. 61:671–683.
- Manooch, C.S. III, and J.C. Potts. 1997c. Age and growth of red snapper, *Lutjanus campechanus*, Lutjanidae, collected along the southeastern United States from North Carolina through the east coast of Florida. J. Elisha Mitchell Scientific Soc. 113:111–122.
- Manooch, C.S. III, J.C. Potts, D.S. Vaughan, and M.L. Burton. 1998. Population assessment of red snapper from the southeastern United States. Fish. Res. 38:19–32.
- Martin, N. 1987. Raw deals. Texas Shores 20:4–8.
- Marx, J.M. 1986. Settlement of spiny lobster, *Panulirus argus*, pueruli in south Florida: an evaluation from two perspectives. Can. J. Fish. Aquat. Sci. 43:2221–2227.
- Matthews, F.D., D.M. Damkaer, L.W. Knapp and B.B. Collette. 1977. Food of western north Atlantic tunas (*Thunnus*) and lancetfishes (*Alepisaurus*). NOAA Tech. Rep. NMFS SSRF-706. 19 p.

- McBride, R.S. 1995. Perennial occurrence and fast growth rates by crevalle jacks (Carangidae: *Caranx hippos*) in the Hudson River estuary. Section VI: pp: 34. In E.A. Blair and J.R. Waldman (Eds.) Final Report of the Tibor T. Polgar Fellowship Program, 1994. Hudson River Foundation, NY.
- McBride, R.S. 2001. Landings, value, and fishing effort for halfbeaks, *Hemiramphus* spp., in the South Florida lampara net fishery. Proc. 52<sup>nd</sup> Ann. Gulf Carib. Fish. Res. Inst:103–115.
- McBride, R.S., and M.D. Murphy. 2003. Current and potential yield per recruit for hogfish, *Lachnolaimus maximus*, in Florida. Proceedings of the 54<sup>th</sup> Annual Meeting of the Gulf and Caribbean Fisheries Institute. Providenciales, Turks and Caicos Islands. 12-17 November 2001. 54:513-525.
- McBride, R.S., and A.K. Richardson. 2007. Evidence of size-selective fishing mortality from an age and growth study of hogfish (Labridae: *Lachnolaimus maximus*), a hermaphroditic reef fish. Bull. Mar. Sci. 80:401-417.
- McBride, R., L. Foushee, and B. Mahmoudi. 1996. Florida's halfbeak, *Hemiramphus* spp., bait fishery. Mar. Fish. Rev. 58:27–36.
- McBride, R.S., T.C. MacDonald, R.E. Matheson, Jr., D.A. Rydene, and P.B. Hood. 2001. Nursery habitats for ladyfish, *Elops saurus*, along salinity gradients in two Florida estuaries. Fish. Bull. 99:443–458.
- McClellan, D.B., and N.J. Cummings. 1997. Preliminary analysis of tag and recapture data of the greater amberjack, *Seriola dumerili*, in the southeastern United States. Proc. 49<sup>th</sup> Gulf Caribb. Fish. Inst. 49:25–45.
- McGovern, J.C., D.M. Wyanski, O. Pashuk, C.S. Manooch II, and G.R. Sedberry. 1998. Changes in the sex ratio and size at maturity of gag, *Mycteroperca microlepis*, from the Atlantic coast of the southeastern United States during 1976–1995. Fish. Bull., U.S.:797–807.
- McGovern, J.C., J.M. Burgos, P.J. Harris, G.R. Sedberry, J.K. Loefer, O. Pashuk, and D. Russ. 2002. Aspects of the life history of red grouper, *Epinephelus morio*, along the southeastern United States. MARFIN Final Report NA97FF0347.
- McInerny, S.A. 2007. Age and growth of red snapper, *Lutjanus campechanus*, from the southeastern United States. Master's Thesis, University of North Carolina Wilmington, Wilmington. 89 p.
- McIntyre, D. K. 2017. Licenses & Fees for Alabama, Florida, Louisiana, Mississippi, and Texas in their Marine Waters for the Year 2016. Gulf States Marine Fisheries Commission. Available from: <http://www.gsmfc.org/publications/GSMFC%20Number%20265.pdf>.
- McMichael, R.H., Jr. 1981. Utilization of the surf zone of a northern Gulf Coast barrier island by the *Menticirrhus* complex (Pisces: Sciaenidae). Master's Thesis, University of Southern Mississippi, Hattiesburg. 86 p.
- McMichael, R.H., Jr., and K.M. Peters. 1989. Early life history of spotted seatrout, *Cynoscion nebulosus* (Pisces: Sciaenidae), in Tampa Bay, Florida. Estuaries 12:98–110.

- McMichael, R.H., Jr., and S.T. Ross. 1987. The relative abundance and feeding habits of juvenile kingfish (Scaenidae: *Menticirrhus*) in a Gulf of Mexico surf zone. *Northeast Gulf Sci.* 9(2):109–123.
- McMillen-Jackson, A. L. & T. M. Bert. 2004. Mitochondrial DNA variation and population genetic structure of the blue crab *Callinectes sapidus* in the eastern United States. *Mar. Biol.* 145:769–777.
- McMillen-Jackson, A.L., T.M. Bert, and P. Steele. 1994. Population genetics of the blue crab *Callinectes sapidus*: modest population structuring in a background of high gene flow. *Mar. Biol.* 118:53–65.
- McRae, G. 1997. 1997 update on Florida's spot fishery. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee.
- Menzel, W. 1989. The biology, fishery, and culture of quahog clams *Mercenaria*. In: Manzi, J.J. and M. Castagna (eds), *Clam Mariculture in North America. Developments in Aquaculture and Fisheries Science*. Vol. 19. Elsevier Publishing Company, New York, pp. 201-242.
- Menzies, R.A., and J.M. Kerrigan. 1979. Implications of spiny lobster recruitment patterns of the Caribbean—a biochemical approach. *Proc. Gulf Caribbean Fisheries Institute* 31:164–178.
- Mercer, L.P. 1989. Fishery management plan for Atlantic croaker (*Micropogonias undulatus*). N.C. Dept. Nat. Res. And Community. Development, Spec. Sci. Rep. 48, 90p.
- Merriman, D. 1940. Morphological and embryological studies on two species of marine catfish, *Bagre marinus* and *Galeichthys felis*. *Zoologica* 25:221–248.
- Merriner, J.V., 1975. Food habits of the weakfish, *Cynoscion regalis*, in North Carolina waters. *Chesapeake Sci.*, 16:74–76.
- Merriner, J. V. 1976. Aspects of the reproductive biology of the weakfish, *Cynoscion regalis* (Scaenidae), in North Carolina. *Fish. Bull., U.S.* 74:18–26.
- Meyer, G.H., and J.S. Franks. 1996. Food of cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. *Gulf Res. Reports* 9:161–167.
- Miles, D.W. 1949. A study of the food habits of fishes of Aransas Bay area. M.S. Thesis, University of Houston, Houston, Texas, 70p
- Miller, R.V. 1969. Continental migrations of fishes—editors note. *Underwat. Nat.* 6:15–23, 44.
- Moe, M.A., Jr. 1969. Biology of the red grouper *Epinephelus morio* (Valenciennes) from the eastern Gulf of Mexico. Professional Papers Ser. 10, Florida Department of Natural Resources Marine Research Laboratory, St. Petersburg, FL.
- Moe, M.A., Jr. 1972. Movement and migration of South Florida fishes. Fl Dept. Nat. Res., Marine Laboratory, Tech. Ser. 69: 25p.

- Mojica, R., Jr., J.M. Shenker, C.W. Harnden, and D.E. Wagner. 1995. Recruitment of bonefish, *Albula vulpes*, around Lee Stocking Island, Bahamas. *Fish. Bull.*, U.S. 93:666–674.
- Moore, J.L. 2001. Age, growth, and reproductive biology of the gray triggerfish (*Balistes capriscus*) from the southeastern United States, 1992-1997. University of Charleston. M.S. Thesis.
- Moss, J., M, D. Behringer, J.D. Shields, A. Baeza, A. Aguilar-Perera, P. G. Bush, C. Dromer, A. Herrera-Moreno, L. Gittens, T. R. Matthews, M.R. McCord, M.T. Schärer, L. Reynal, N. Truelove, M.J. Butler. 2013. Distribution, prevalence, and genetic analysis of *Panulirus argus* virus 1 (PaV1) from the Caribbean Sea. *Diseases of Aquatic Organisms* 104:129-140. doi: 10.3354/dao02589.
- Muller, R.G., D. Chagaris, T.M. Bert, C. Crawford, and Ryan Gandy. 2011. The 2011 stock assessment update for the stone crab, *Menippe* spp., fishery in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2011-003.
- Muller, R.G., and M.D. Murphy. 1994. A stock assessment of sheepshead. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1994-005)
- Muller, R.G., and R. G. Taylor. 2006. The 2005 stock assessment update of common snook, *Centropomus undecimalis*. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2006-XXX.
- Muller, R.G., and R.G. Taylor. 2012. The 2012 stock assessment update of common snook, *Centropomus undecimalis*. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2012-002.
- Muller, R.G., and R.G. Taylor. 2013. The 2013 stock assessment update of common snook, *Centropomus undecimalis*. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2013-004.
- Muller, R.G., T. M. Bert, and S.D. Gerhart 2006. The 2006 update for the stone crab, *Menippe* spp., fishery in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2006-011: 47p.
- Muller, R.G., J.H. Hunt, T.R. Matthews, and W.C. Sharp. 1997. Evaluation of effort reduction in the Florida Keys spiny lobster, *Panulirus argus*, fishery using age-structured population analysis. *Mar. Freshwater Res.* 48:1045–1058.
- Muller, R.G., M.D. Murphy, J.A. De Silva, L.R. Barbieri. 2003. A stock assessment of yellowtail snapper, *Ocyurus chrysurus*, in the Southeast United States. Final Report Submitted to the National Marine Fisheries Service, the Gulf of Mexico Fishery Management Council, and the South Atlantic Fishery Management Council as part of Southeast Data, Assessment, and Review (SEDAR) 03. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2003-10.

- Muller, R.G., D. Chagaris, T.M. Bert, C. Crawford, and R. Gandy. 2011. The 2011 stock assessment update for the stone crab, *Menippe spp.*, fishery in Florida. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2011-003.
- Muller, R.G., W.C. Sharp, T.R. Mathews, R. Bertelsen, and J.H. Hunt. 1999. The 1999 update of the stock assessment for the Florida Keys spiny lobster, *Panulirus argus*. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1999-009)
- Muller, R.G., W.C. Sharp, T.R. Mathews, R. Bertelsen, and J.H. Hunt. 2000. The 2000 update of the stock assessment for spiny lobster, *Panulirus argus*, in the Florida Keys. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2000-007)
- Muller, R.G., A.A. Trotter, and P.W. Stevens. 2015. The 2015 stock assessment update of Common Snook, *Centropomus undecimalis*. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2015-004.
- Muncy, R.J., and W. M. Wingo. 1983. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico)—sea catfish and gafftopsail catfish. U.S. Fish and Wildl. Serv., Div Biological Serv., FWS/OBS-82/11.5. U.S. Army Corps of Engineers, TR EL-82-4.
- Munyandorero, J., M. Masi, and S. D. Allen. 2017. An assessment of the status of sheepshead in Florida waters through 2015. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2017-002.
- Munyandorero, J., B. Mahmoudi, and C. Bradshaw. 2007. An assessment of the status of silver mullet in Florida waters through 2006. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2007-XXX.
- Munyandorero, J., M.D. Murphy, and T.C. MacDonald. 2006. An assessment of the status of sheepshead in Florida waters through 2004. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2006-009.
- Murie, D.J., and D.C. Parkyn. 2002. Comparison of total mortality of white grunt from the head-boat fishery on the Gulf coast of Florida during spawning and postspawning seasons. N. Am. J. Fish. Manage. 22: 806-814.
- Murie, D.J., and D.C. Parkyn. 2005. Age and growth of white grunt (*Haemulon plumierii*): a comparison of two populations along the Florida west coast. Bull. Mar. Sci. 76(1):73-93.
- Murphy, M.D. 2002. A stock assessment of red drum, *Sciaenops ocellatus*, in Florida: status of the stocks through 2000. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2006-016)

- Murphy, M.D. 2005. A stock assessment of red drum, *Sciaenops ocellatus*, in Florida: status of the stocks through 2003. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2005-006.
- Murphy, M.D. 2010. Regional analysis of Florida's Gulf and Atlantic stocks of Red Drum. Report to the Florida FWCC Division of Marine Fisheries Management from the Fish and Wildlife Research Institute. August, 2010.
- Murphy, M.D., and R.E. Crabtree. 2001. Changes in the age structure of nearshore adult red drum off west-central Florida related to recruitment and fishing mortality. *N. Am. J. Fish. Manage.* 21:671–678.
- Murphy, M.D., and T.C. MacDonald. 2000. An assessment of the status of sheepshead in Florida waters through 1999. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2000-009)
- Murphy, M.D., and R.G. Muller. 1995. A stock assessment of black drum *Pogonias cromis* in Florida. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1995-005:54p.)
- Murphy, M.D., and R.G. Taylor. 1989. Reproduction and growth of black drum, *Pogonias cromis*, in northeast Florida. *Northeast Gulf Sci.* 10:127–137.
- Murphy, M.D. and R.G. Taylor. 1990. Reproduction, growth, and mortality of red drum *Sciaenops ocellatus*, in Florida waters. *Fish. Bull.* 88 (3), 531–542.
- Murphy, M.D., and R.G. Taylor. 1994. Age, growth, and mortality of spotted seatrout in Florida waters. *Trans. Amer. Fish. Soc.* 123:482–497.
- Murphy, M.D., D. Chagaris, and D.T. Addis. 2011. An assessment of the status of spotted seatrout in Florida waters through 2009. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2011-002.
- Murphy, M.D., C.B. Guenther, and B. Mahmoudi. 2006. An assessment of the status of spotted seatrout in Florida waters through 2005. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2006-XXX:195p.
- Murphy, M.D., B. Mahmoudi, R.G. Muller. 1997. A stock assessment of sheepshead. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1997-004)

- Murphy, M.D., A.L. McMillen-Jackson, and B. Mahmoudi. 2007. A stock assessment for blue crab, *Callinectes sapidus*, in Florida waters. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2007-006.
- Murphy, M.D., C.A. Meyer, and A.L. McMillen-Jackson. 2001. A stock assessment for blue crab, *Callinectes sapidus*, in Florida waters. Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR2001-008)
- Murphy, M.D., R.G. Muller, and P.B. Hood. 1996. A stock assessment of Florida pompano *Trachinotus carolinus*. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1996-XXX)
- Murphy, M.D., R.G. Muller, and K. Guindon. 2008. A stock assessment for pompano, *Trachinotus carolinus*, in Florida waters through 2005. Florida Fish and Wildlife Conservation Commission. Fish and Wildlife Research Institute In House Report IHR2008-004.
- Murphy, M.D., R.G. Muller, and B. McLaughlin. 1994. A stock assessment of southern and gulf flounder. Florida Marine Research Institute report to the Florida Marine Fisheries Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1994-003)
- Murphy, M.D., D.J. Murie, and R.G. Muller. 1999. Stock assessment of white grunt from the west coast of Florida. Florida Marine Research Institute report to the Florida Fish and Wildlife Conservation Commission, Tallahassee. (currently Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute In-House Report IHR1999-007)
- Music, J.L. and J. Pafford. 1984. Population dynamics and life history aspects of major marine sportfishes in Georgia's coastal waters. Georgia Dept. Nat. Res., Coastal res. Div. Contr. Ser. Num. 38. 382 pp.
- Myer, G.H. and J.S. Franks. 1996. Food of cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. Gulf Res. Reports 9:161–167.
- Nance, J.M., E.F. Klima, and T.E. Czaplá. 1989. Gulf of Mexico shrimp stock assessment workshop. NOAA Tech. Memo. NMFS-SEFC-239.
- Nance, J. M. 1999. Stock assessment report 1998, Gulf of Mexico shrimp fishery. NMFS SEFSC Galveston Laboratory, Galveston, Texas. Unpublished Document, 13 pp.
- Nance, J.M. 2007. Review of the status and health of the shrimp stocks for 2006. Report to the Gulf of Mexico Fisheries Management Council, Tab D, No. 3, 2007 meeting.
- Nash, C.E., C. M. Kuo, and S.C. McConnel. 1974. Operational procedures for rearing larvae of the grey mullet (*Mugil cephalus* Linnaeus). Aquaculture 3:15–24.

National Marine Fisheries Service Southeast Fisheries Science Center staff. 2002 (Draft December 14, 2002). Status of red grouper in United States waters of the Gulf of Mexico during 1986-2001. NMFS-SEFSC Sustainable Fisheries Division Contribution No. SFD-01/02-175rev. 65p.

Naughton, S.P. and C.H. Saloman. 1984. Food of bluefish (*Pomatomus saltatrix*) from the U.S. south Atlantic and Gulf of Mexico. NOAA Tech. Mem. NMFS-SEFC-150.

Naughton, S.P. and C.H. Saloman. 1985. Food of gag (*Mycteroperca microlepis*) from North Carolina and three areas of Florida. NOAA Tech. Mem. NMFS-SEFC-160.

NCDMF 2015. Stock Assessment of Spotted Seatrout, *Cynoscion nebulosus*, in Virginia and North Carolina Waters 2014. NCDMF SAP-SAR-2015-02.

Nelson, G.A. 1998. Abundance, growth, and mortality of young-of-the-year pinfish, *Lagodon rhomboides*, in three estuaries along the gulf coast of Florida. Fish. Bull. 96:315–328.

Nelson, G.A. 2002. Age, growth, mortality, and distribution of pinfish (*Lagodon rhomboides*) in Tampa Bay and adjacent Gulf of Mexico waters. Fish. Bull. 100:582–592.

Nelson, R.S., and C.S. Manooch III. 1982. Growth and mortality of red snappers in the west-central Atlantic Ocean and northern Gulf of Mexico. Trans. Amer. Fish. Soc. 111:465–475.

Odum, W.E. 1970. Utilization of the direct grazing and plant detritus food chains by the striped mullet, *Mugil cephalus*. p. 222–240. In: J.J. Steele (editor), *Marine Food Chains*, Oliver and Boyd, Ltd., Edinburgh, Scotland.

Odum, W.E. 1971. Pathway of energy flow in a South Florida estuary. Univ. Miami Sea Grant Tech. Bull. 7, 160p.

Ogburn, M.V. 1984. Feeding ecology and the role of algae in the diet of sheepshead *Archosargus probatocephalus* (Pisces: Sparidae) on two North Carolina jetties. M.S. Thesis, Univ. North Carolina at Wilmington.

O’Hop, J., M. Murphy, and D. Chagaris. 2012. The 2012 stock assessment report for yellowtail snapper in the South Atlantic and Gulf of Mexico. Florida Fish and Wildlife Conservation Commission. Fish and Wildlife Research Institute. 63p.

Olla, B.L., H.M. Katz, and A.L. Studholme. 1970. Prey capture and feeding motivation in the bluefish, *Pomatomus saltatrix*. Copeia 1970(2):360–362.

Overstreet, R. M. and R. W. Heard 1982. Food contents of six commercial fishes from Mississippi Sound. Gulf Res. Reports 7(2):137–149.

Oxenford, H.A. and W. Hunte. 1983. Age and growth of dolphin, *Coryphaena hippurus*, as determined by growth rings from otoliths. Fish. Bull., U.S. 81:906–909.

- Oxenford, H.A., and W. Hunte. 1986. A preliminary investigation of the stock structure of the dolphin, *Coryphaena hippurus*, in the western central Atlantic. *Fish Bull.* 84:451–469.
- Oxenford, H.A. and W. Hunte. 1999. Feeding habits of the dolphin (*Coryphaena hippurus*) in the eastern Caribbean. *Scientia Marina* 63 (2–3).
- Pacheco, A.L. 1962. Age and growth of spot in lower Chesapeake Bay, with notes on distribution and abundance of juveniles in the York River system. *Chesapeake Sci.* 3:18–28.
- Palko, B.J. 1984. An evaluation of hard parts for age determination of pompano (*Trachinotus carolinus*), ladyfish (*Elops saurus*), crevalle jack (*Caranx hippos*), gulf flounder (*Paralichthys albigutta*), and southern flounder (*Paralichthys lethostigma*). NOAA Tech. Memo. NMFS-SEFC-132, 11p.
- Palko, B.J., G.L. Beardsley, and W.J. Richards. 1982. Synopsis of the biological data on dolphin-fishes, *Coryphaena hippurus* (Linnaeus) and *Coryphaena equiselis* (Linnaeus). NOAA Tech Report NMFS Circular 443.
- Parker, J.C. 1971. The biology of the spot, *Leiostomus xanthurus* Lacepede, and Atlantic croaker, *Micropogon undulatus* (Linnaeus) in two Gulf of Mexico nursery areas. Ph.D. Diss. Texas A&M University, College Station.
- Pearson, J., 1929. Natural history and conservation of redfish and other commercial Sciaenidae of Texas. *Bull. U.S. Bur. Fish. Wash.* 44:129–214.
- Perez, A. and R. Rubio. 1986. Análisis electrónico de frecuencias de largo en poblaciones de peces cubanos. 5to. Foro Cient. Cip, Cent. Invest. Pesq. Cuba, En. 1986. [as cited in FISHBASE]
- Peters, J.S., and D.J. Schmidt. 1997. Daily age and growth of larval and early juvenile Spanish mackerel, *Scomberomorus maculatus*, from the South Atlantic Bight. *Fish. Bull., U.S.* 95:530–539.
- Peters, K.M., R.E. Matheson, Jr., and R.G. Taylor. 1998. Reproduction and early life history of common snook, *Centropomus undecimalis* (Bloch), in Florida. *Bull. Mar. Sci.* 62:509–529.
- Peters, K.M., and R.G. McMichael, Jr. 1987. Early life history of *Sciaenops ocellatus* (Pisces: Sciaenidae) in Tampa Bay, Florida. *Estuaries* 10:92–107.
- Phares, P.L. 1992. Analysis of trends in catch, effort, and catch per unit effort in the Florida stone crab (genus *Menippe*) fishery, p. 18–25 *In*: T.M. Bert (editor), Proceedings of a symposium on stone crab (genus *Menippe*) biology and fisheries. Florida Mar. Res. Pub. 50.
- Piedra, G. 1969. Materials on the biology of the yellowtail snapper (*Ocyurus chrysurus* Black). Pages 251–296 *in* A.S. Bogdanov, editor. Soviet-Cuban Fishery Research. Translated from Russian: Israel Program for Scientific Translations, Jerusalem, Israel.
- Porch, C.E. 2000. Status of the red drum stocks in the Gulf of Mexico. Version 2.1. Report from the NMFS-SEFSC Sustainable Fisheries Division, Miami, Contribution: SFD-98/99-76.

- Porch, C.E., C.A. Wilson, and D.L. Nieland. 2002. A new growth model for Red Drum (*Sciaenops ocellatus*) that accommodates seasonal and ontogenic changes in growth rates. *Fishery Bulletin* 100:149-152.
- Pottern, G.B., M.T. Huish, and J.H. Kerby. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Mid-Atlantic)—bluefish. U.S. Fish Wildl. Serv. Biol. Rep. 82 (11.94). U.S. Army Corps of Engineers. TR EL-82-4. 20p.
- Potts, J.C., 2000. Population assessment of two stocks of white grunt, *Haemulon plumieri*, from the Southeastern coast of the United States. NOAA report prepared for the South Atlantic Fishery Management Council. 66 pp.
- Potts, J.C. and K. Brennan. 2001. Trends in catch data and estimated static SPR values for fifteen species of reef fish landed along the Southeastern United States. National Oceanic and Atmospheric Administration, Center for Coastal Fisheries and Habitat Research. Beaufort Laboratory. Beaufort, NC. February 2001.
- Potts, J.C., and C.S. Manooch III. 2001. Differences in age and growth of white grunt from North Carolina and South Carolina compared with southeast Florida. *Bull. Mar. Sci.* 68:1–12.
- Potts, J.C., C.S. Manooch III, and D.S. Vaughn. 1998. Age and growth of vermilion snapper from the Southeastern United States. *Trans. Amer. Fish. Soc.* 127:787–795.
- Powell, D. 1975. Age, growth, and reproduction in Florida stocks of Spanish mackerel, *Scomberomorus maculatus*. *Fla. Mar. Res. Publ.* 5:21p.
- Prager, M.H. 2000. Exploratory assessment of the dolphin, *Coryphaena hippurus*, based on the U.S. landings from the Atlantic Ocean and Gulf of Mexico. Report from the SEFSC Beaufort Laboratories, NOAA-NMFS to the South Atlantic Fishery Management Council, Charleston, South Carolina.
- Prest, K.W. 1971. Fundamentals of sexual maturation, spawning, and fecundity of the thread herring (*Opisthonema oglinum*) in the eastern Gulf of Mexico. NMFS Biol. Lab, St. Petersburg Beach, Florida. Unpublished manuscript.
- Ramsey, P.R., and J.M. Wakeman. 1987. Population structure of *Sciaenops ocellatus* and *Cynoscion nebulosus* (Pisces: Sciaenidae): biochemical variation, genetic subdivision and dispersal. *Copeia* 1987:682–695.
- Randall, J.E. 1967. Food habits of reef fishes of the West Indies. *Stud. Trop. Ocean.* Miami 5:665–847.
- Reintjes, J.W. 1979. Coastal herrings and associated species: a profile of species or groups of species, their biology, ecology, current exploitation with comments on economic and social information. NMFS-SEFSC Beaufort, N.C. 170pp.

- Render, J.H., and C.A. Wilson. 1992. Reproductive biology of sheepshead in the northern Gulf of Mexico. *Trans. Amer. Fish. Soc.* 121:757–764.
- Restrepo, V.R, G.G. Thompson, P.M. Mace, W.L. Gabriel, L.L. Low, A.D. MacCall, R.D. Methot, J.E. Powers, B.L. Taylor, P.R. Wade, and J.F. Witzig. 1998. Technical guidance on the use of precautionary approaches to implementing National Standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. DOC, NOAA, NMFS, NOAA Technical Memorandum NMFS-F/SPO-31. 54 pp.
- Richards, C.E. 1967. Age, growth, and fecundity of the cobia, *Rachycentron canadum*, from the Chesapeake Bay and adjacent mid-Atlantic waters. *Trans. Amer. Fish. Soc.* 96:343–350.
- Richardson, L.R., and J.R. Gold. 1997. Mitochondrial DNA diversity in and population structure of red grouper, *Epinephelus morio*, from the Gulf of Mexico. *Fish. Bull., U.S.* 95:174–179.
- Rickards, W.L. 1968. Ecology and growth of juvenile tarpon, *Megalops atlanticus*, in a Georgia salt marsh. *Bull. Mar. Sci.* 18:220–239.
- Rivera, G.A. and R.S. Appeldoorn. 2000. Age and growth of dolphin, *Coryphaena hippurus*, off Puerto Rico. *Fish. Bull., U.S.* 98:345–352.
- Robillard E, Reiss CS, Jones CM. 2008. Reproductive biology of bluefish (*Pomatomus saltatrix*) along the East Coast of the United States. *Fisheries Research* 90 (2008): 198-208.
- Robillard, E., C.S. Reiss, and C.M. Jones. 2009. Age-validation and growth of bluefish (*Pomatomus saltatrix*) along the East Coast of the United States. *Fisheries Research* 95:65-75.
- Robins, C.R. 1978. The tarpon – unusual biology and man’s impact determine its future. *Mar. Rec. Fish.* 2:105–112.
- Roelke, L.A., and L.A. Cifuentes. 1997. Use of stable isotopes to assess groups of king mackerel, *Scomberomorus cavalla*, in the Gulf of Mexico and southeastern Florida. *Fish. Bull., U.S.* 95:540–551.
- Rooker, J.R., S.A. Holt, G.J. Holt, and L.A. Fuiman. 1999. Spatial and temporal variability in growth, mortality, and recruitment potential of postsettlement red drum, *Sciaenops ocellatus*, in a subtropical estuary. *Fish. Bull., U.S.* 97:581–590.
- Rooker, J.R., G.W. Stunz, S.A. Holt, and T.J. Minello. 2010. Population connectivity of red drum in the northern Gulf of Mexico. *Mar. Ecol. Prog. Ser.* 407:187-196.
- Rose, C.D., and W.W. Hassler. 1968. Age and growth of the dolphin, *Coryphaena hippurus* (Linnaeus), in North Carolina waters. *Trans. Amer. Fish. Soc.* 97:271–276.
- Ross, J.L., T.M. Stevens, and D.S. Vaughan. 1995. Age, growth, mortality, and reproductive biology of red drums in North Carolina waters. *Trans. Amer. Fish. Soc.* 124:37–54.

- Ross, S.W. 1988. Age, growth, and mortality of Atlantic croaker in North Carolina, with comments on population dynamics. *Trans. Amer. Fish. Soc.* 117:461–473.
- Rothschild, B.J. and J.S. Ault. 1992. Assessment of the Chesapeake Bay Blue Crab Stock. Ref.No. UMCEES[CBL] 92-082. Final Report for NO/NMFS Grant NA16FU0529-01. 201p.
- Rozas, L.P., and C.T. Hackney. 1984. Use of oligohaline marshes by fishes and macrofaunal crustaceans in North Carolina. *Estuaries* 7:213–224.
- Rugolo, L., and eight other authors. 1997. Stock assessment of Chesapeake Bay blue crab (*Callinectes sapidus*). Report by the Technical Subcommittee of the Chesapeake Bay Stock Assessment Committee of the National Marine Fisheries Service, NOAA, April 1997.
- Rutherford, E.S., T.W. Schmidt, and T.J. Tilmant. 1989a. Population structure, food habits, and spawning activity of gray snapper, *Lutjanus griseus*, in Everglades National Park. So. Fla. Res. Center Rep. SFRC-83/02. U.S. National Park Service, Homestead, FL.
- Rutherford, E.S., J.T. Tilmant, E.B. Thue, and T.W. Schmidt. 1989b. Fishery harvest and population dynamics of gray snapper, *Lutjanus griseus*, in Florida Bay and adjacent waters. *Bull. Mar. Sci.* 44:139–154.
- Saloman, C.H., and S.P. Naughton. 1983a. Food of king mackerel, *Scomberomorus cavalla*, from the southeastern United States including the Gulf of Mexico. U.S. Dept. Commerce NOAA Tech. Memo. NMFS-SEFC-126. 25p.
- Saloman, C.H., and S.P. Naughton. 1983b. Food of Spanish mackerel, *Scomberomorus maculatus*, from the Gulf of Mexico and southeastern seaboard of the United States. U.S. Dept. Commerce NOAA Tech. Memo. NMFS-SEFC-128. 22p.
- Saloman, C. H., and S. P. Naughton. 1984. Food of crevalle jack (*Caranx hippos*) from Florida, Louisiana, and Texas. U.S. Dep. Commerce, NOAA Tech. Memo. NMFS-SEFC-134, 34 p.
- Saloman, C.H., and J.L. Taylor. 1969. Age and growth of large southern quahogs from a Florida estuary. *Proc. Natl. Shellfish Assoc.* 59:46–51.
- Schaaf, W. E. 1979. An analysis of the dynamic population response of Atlantic menhaden, *Brevoortia tyrannus*, to an intensive fishery. *Rapp. P.V. Reun. Cons. Int. Explor. Mer.* 177:243–251.
- Schaefer, R.H. 1965. Age and growth of the northern kingfish in New York waters. *N.Y. Fish Game J.* 12:191–216.
- Scharf, F.S., J.P. Manderson, M.C. Fabrizio, J.P. Pessutti, J.E. Rosendale, R.J. Chant, and A.J. Bejda. 2004. Seasonal and interannual patterns of distribution and diet of bluefish within a middle Atlantic Bight estuary in relation to abiotic and biotic factors. *Estuaries.* 27:426-436.

- Schirripa, M.J. 1998. Status of the vermilion snapper fishery of the Gulf of Mexico: assessment 4.0. NMFS-SEFSC Sustainable Fisheries Division Contr. No. SFD-97/98-09.
- Schirripa, M.J. and K.M. Burns. 1997. Growth estimates for three species of reef fish in the eastern Gulf of Mexico. Bull Mar. Sci. 61:581–591.
- Schirripa, M.J., and C.P. Goodyear. 1994. Status of the gag stocks of the Gulf of Mexico. Assessment 1.0. Report from the NMFS SEFSC Sustainable Fisheries Division Miami.
- Schirripa, M.J., and C.M. Legault. 1997. Status of the red snapper stocks of the Gulf of Mexico: updated through 1996. Report from the NMFS-SEFSC Sustainable Fisheries Division, Miami, Contribution: MIA-97/98-05.
- Schirripa, M.J., and C.M. Legault. 1999. Status of the red snapper stocks of the Gulf of Mexico: updated through 1998. Report from the NMFS-SEFSC Sustainable Fisheries Division, Miami, Contribution: SFD-99/00-75.
- Schirripa, M.J., C.M. Legault, and M. Ortiz. 1999. The red grouper fishery of the Gulf of Mexico. Assessment 3.0. Report from the NMFS-SEFSC Sustainable Fisheries Division, Miami, Contribution: SFD-98/99-56.
- Schmidt, D.J., M.R. Collins, and D.M. Wyanski. 1993. Age, growth, maturity, and spawning of Spanish mackerel, *Scomberomorus maculatus* (Mitchill), from the Atlantic coast of the southeastern United States. Fish. Bull. 91:526–533.
- Schwartz, F.J., and H.J. Porter. 1977. Fishes, macroinvertebrates, and their ecological inter-relationships with a calico scallop bed off North Carolina. Fish. Bull. U.S. 75:427–446.
- SEDAR 1. 2002. Stock assessment of south Atlantic red porgy. South Atlantic Fishery Management Council SEDAR1 Assessment Report, Charleston, S.C. 102p.
- SEDAR 5 and South Atlantic Fishery Management Council. 2004a. Complete stock assessment report of SEDAR 5: Atlantic and Gulf of Mexico king mackerel. South Atlantic Fishery Management Council SEDAR5 Assessment Reports 1-5, Charleston, S.C. 205p.
- SEDAR 6 and South Atlantic Fishery Management Council. 2004b. Complete stock assessment report of SEDAR 6: hogfish snapper. South Atlantic Fishery Management Council SEDAR6 Assessment Report 2, Charleston, S.C. 127p.
- SEDAR 07. 2005. Assessment Summary Report, Gulf of Mexico red snapper. Nat. Mar. Fish. Serv., SEDAR, Charleston, S.C. 13p.
- SEDAR 08 U.S. Stock Assessment Panel. 2005. Assessment of spiny lobster, *Panulirus argus*, in the southeast United States. National Marine Fisheries Service SEDAR 08 Stock assessment report, Charleston, S.C., 29 April 2005, 97p.

SEDAR 08 Update Stock Assessment Panel. 2010. Update Spiny Lobster Stock Assessment Workshop Report, Key West, FL., 122p.

SEDAR 09 Stock Assessment Panel. 2006. Greater amberjack, *Seriola dumerili*, in the Gulf of Mexico stock assessment report. National Marine Fisheries Service, SEDAR, Charleston, S.C. 61 p.

SEDAR 09. 2006. Stock assessment report of SEDAR 9: Gulf of Mexico gray triggerfish. SEDAR 9 Assessment Report 1. National Marine Fisheries Service, SEDAR, Charleston, S.C. 61p.

SEDAR 09. 2006b. Stock assessment report of SEDAR 9: Gulf of Mexico vermilion snapper. SEDAR 9 Assessment Report 3. National Marine Fisheries Service, SEDAR, Charleston, S.C. 61p.

SEDAR 09 Update. 2011a. SEDAR 9 stock assessment update report: Gulf of Mexico greater amberjack. National Marine Fisheries Service, SEDAR, Charleston, S.C. 179p.

SEDAR 09 Update. 2011b. SEDAR 9 stock assessment update report: Gulf of gray triggerfish. National Marine Fisheries Service, SEDAR, Tampa, FL. 261p.

SEDAR 09 Update. 2011c. SEDAR 9 stock assessment update report: Vermilion snapper. National Marine Fisheries Service, SEDAR, Tampa, FL. 87p.

SEDAR 10 Review Panel. 2006a. SEDAR 10 review workshop report: South Atlantic gag grouper. National Marine Fisheries Service, Southeast Fisheries Science Center SEDAR, June 26-30, 2006, Atlanta, Georgia

SEDAR 10 Review Panel. 2006b. SEDAR 10 review workshop report: Gulf of Mexico gag grouper. National Marine Fisheries Service, Southeast Fisheries Science Center SEDAR, June 26-30, 2006, Atlanta, Georgia.

SEDAR 10 Assessment Workshop Panel. 2006. SEDAR 10 stock assessment report 2: Gulf of Mexico gag grouper. National Marine Fisheries Service, Southeast Fisheries Science Center SEDAR. May 1-5, 2006, Miami, Florida.

SEDAR 12. 2006. Stock assessment report of SEDAR 12: Gulf of Mexico red grouper. SEDAR 12 Assessment Report 1. National Marine Fisheries Service, SEDAR, Charleston, S.C. 358p.

SEDAR 15. 2008a. Stock assessment report of SEDAR 15: South Atlantic red snapper. SEDAR 15 Assessment Report 1. South Atlantic Fishery Management Council, SEDAR, February 2008, North Charleston, S.C. 212p.

SEDAR 15. 2008b. Stock assessment report of SEDAR 15: South Atlantic greater amberjack. SEDAR 15 Assessment Report 2. South Atlantic Fishery Management Council, SEDAR, February 2008, North Charleston, S.C. 379p.

- SEDAR 16. 2009. Stock assessment report of SEDAR 16: South Atlantic and Gulf of Mexico king mackerel. SEDAR 16 Assessment Report 1. South Atlantic Fishery Management Council, SEDAR, March 2009, North Charleston, S.C. 484p.
- SEDAR 17. 2008a. Stock assessment report of SEDAR 17: South Atlantic spanish mackerel. SEDAR 17 Assessment Report 1. South Atlantic Fishery Management Council, SEDAR, November 2008, North Charleston, S.C. 508p.
- SEDAR 17. 2008b. Stock assessment report of SEDAR 17: South Atlantic vermilion snapper. SEDAR 17 Assessment Report 2. South Atlantic Fishery Management Council, SEDAR, November 2008, North Charleston, S.C. 450p.
- SEDAR 17. 2012. Stock assessment of vermilion snapper of the Southeastern United States, SEDAR update assessment. Southeast Fisheries Science Center, National Marine Fisheries Service, October 2012, Miami, FL. 110p.
- SEDAR 18. 2009. Stock assessment report of SEDAR 18: Atlantic red drum. SEDAR 18 Assessment Report 1. South Atlantic Fishery Management Council, SEDAR, October 2009, North Charleston, S.C. 56p.
- SEDAR 19. 2010. Stock assessment report of SEDAR 19: South Atlantic red grouper. SEDAR 19 Assessment Report 2. South Atlantic Fishery Management Council, SEDAR, April 2010, North Charleston, S.C. 612p.
- SEDAR 20 and Atlantic States Marine Fisheries Commission. 2010a. Atlantic menhaden stock assessment and review panel reports. SEDAR 20 Assessment Report 1. ASMFC Stock assessment report no. 10-02. Atlantic States Marine Fisheries Commission, Washington, D.C. 328p.
- SEDAR 20 and Atlantic States Marine Fisheries Commission. 2010b. Atlantic croaker 2010 stock assessment report draft for peer review. SEDAR 20 Assessment Report 2. Atlantic States Marine Fisheries Commission, Washington, D.C. 328p.
- SEDAR. 2012. The 2012 stock assessment report for Yellowtail Snapper in the South Atlantic and Gulf of Mexico. SEDAR, North Charleston, SC. 257pp. available online at: <http://sedarweb.org/sedar-27a>
- SEDAR 27. 2011. Stock assessment report of SEDAR 27: Gulf of Mexico Menhaden. D. Vaughan, A. Schueller, J. Smith, and S. VanderKooy (eds.). North Charleston, S.C. 492 p.
- SEDAR 27A. 2012. The 2012 Stock assessment report for Yellowtail Snapper in the South Atlantic and Gulf of Mexico. J. O'Hop, M. Murphy, and D. Chagaris. St. Petersburg, FL. 341 pp.
- SEDAR 31. 2013. SEDAR 31 – Gulf of Mexico Red Snapper Stock Assessment Report. SEDAR, North Charleston SC. 1103 pp.
- SEDAR 33. 2014a. SEDAR 33 – Gulf of Mexico Gag Stock Assessment Report. SEDAR, North Charleston SC. 609 pp.

- SEDAR 33. 2014b. SEDAR 33 – Gulf of Mexico Greater Amberjack Stock Assessment Report. SEDAR, North Charleston SC. 490 pp.
- SEDAR 33. 2016. SEDAR 33 – UPDATE REPORT Gulf of Mexico Gag Grouper. SEDAR, North Charleston SC. 128 pp.
- SEDAR 38. 2014a. SEDAR 38 – Gulf of Mexico King Mackerel Stock Assessment Report. SEDAR, North Charleston, SC. 465 pp.
- SEDAR 38. 2014b. SEDAR 38 – South Atlantic King Mackerel Stock Assessment Report. SEDAR, North Charleston, SC. 502 pp.
- SEDAR 40. 2015. SEDAR 40 – Atlantic Menhaden Stock Assessment Report. SEDAR, North Charleston, SC. 643 pp.
- SEDAR 41. 2016. SEDAR 41 – South Atlantic Red Snapper and Gray Triggerfish Assessment Report. SEDAR, North Charleston SC. 660 pp. available online at: <http://sedarweb.org/sedar-41>.
- SEDAR 42. 2015. SEDAR 42 – Gulf of Mexico Red Grouper: SECTION III: Assessment Process Report. SEDAR, North Charleston, SC. 612 pp.
- SEDAR 43. 2015. SEDAR 43 – Gulf of Mexico Gray Triggerfish. SEDAR, North Charleston, SC. 193 pp. available online at: [http://sedarweb.org/docs/sar/S43\\_SAR\\_FINAL.pdf](http://sedarweb.org/docs/sar/S43_SAR_FINAL.pdf).
- SEDAR 44. 2015. SEDAR 44 – Atlantic Red Drum Stock Assessment Report. SEDAR, North Charleston, SC. 890 pp.
- SEDAR 45. 2016. SEDAR 45 – Gulf of Mexico Vermilion Snapper Stock Assessment Report. SEDAR. North Charleston, SC. 188 pp.
- SEDAR 49. 2016. SEDAR 49 – Gulf of Mexico Data-limited Species: Red Drum, Lane Snapper, Wenchman, Yellowmouth Grouper, Speckled Hind, Snowy Grouper, Almaco Jack, Lesser Amberjack. SEDAR. North Charleston, SC. 618 pp.
- SEDAR 51. 2018. SEDAR 51 – Gulf of Mexico Gray Snapper. SEDAR. North Charleston, SC. 428 pp.
- SEDAR 53. 2017. SEDAR 53 – South Atlantic Red Grouper Assessment Report. SEDAR, North Charleston, SC. 159 pp. available online at: <http://sedarweb.org/sedar-53>.
- SEDAR 55. 2018. SEDAR 55 – South Atlantic Vermilion Snapper Assessment Report. SEDAR, North Charleston, SC. 170 pp. available online at: <http://sedarweb.org/sedar-55>.
- SEDAR 57. 2019. SEDAR 57 – U.S. Caribbean Spiny Lobster Assessment Report. SEDAR, North Charleston, SC. 232 pp. available online at: <http://sedarweb.org/sedar-57>.

- Sekavec, G.B. 1974. Summer foods, length-weight relationship, and condition factor of juvenile ladyfish, *Elops saurus* Linnaeus, from Louisiana coastal streams. *Trans. Amer. Fish. Soc.* 103:472–476.
- Seyoum, S., M.D. Tringali, B.L. Barthel, V. Villanova, C. Puchulutegui, M.C. Davis, A.C. Alvarez. 2014. Stock Boundaries for Spotted Seatrout (*Cynoscion nebulosus*) in Florida Based on Population Genetic Structure. Florida Fish and Wildlife Research Institute Technical Reports TR-18.
- Seyoum, S., M.D. Tringali, T.M Bert, D. McElroy, R. Stokes. 2000. An analysis of genetic population structure in red drum, *Sciaenops ocellatus*, based on mtDNA control region sequences. *Fish. Bull., U.S.* 98:127–138.
- Seyoum, S., R. S. McBride, C. Puchutulegui, J. Dutka-Gianelli, A. C. Alvarez, & K. Panzner. 2017. Genetic population structure of a coastal marine fish (Sheepshead; *Archosargus probatocephalus* [Sparidae]) in the southeastern United States: multiple population clusters based on species-specific microsatellite markers. *Journal of Marine Biology* (in press).
- Seyoum, S., R.S. McBride, M.D. Tringali, V.L. Villanova, C. Puchutulegui, S. Gray, N. Van Bibber. 2018. Genetic population structure of the spotted seatrout (*Cynoscion nebulosus*): simultaneous examination of the mtDNA control region and microsatellite marker results. *Bull. Mar. Sci.* 94:1-25.
- Shaffer, R.V., and E.L. Nakamura. 1989. Synopsis of biological data on the cobia *Rachycentron canadum* (Pisces: Rachycentridae). NOAA Tech. Rep. NMFS 82.
- Sharp, W. C., R.D. Bertelsen, and V.R. Leeworthy. 2005 Long-term trends in the recreational lobster fishery of Florida, United States: landings, effort and implications for management. *New Zealand Jour. of Mar. and Freshwat. Res.* 39: 733-747.
- Sharp, W.C., W.A. Lellis, M.J. Butler, W.F. Herrnkind, J.H. Hunt, M. Pardee-Woodring and T.R. Matthews. 2000. The Use of Coded Microwire Tags in Mark-Recapture Studies of Juvenile Caribbean Spiny Lobster, *Panulirus argus*. *Journal of Crustacean Biology.* 20: 510-521.
- Sharp, W., N. Sheridan. 2019. Trend Analysis of Marine Life Landings 2005-2017. Report to the Florida Fish and Wildlife Conservation Commission's Division of Marine Fisheries Management.
- Shepard, K.E., W.F. Patterson, and D.A. DeVries. 2010. Trends in Atlantic contribution to mixed-stock king mackerel landings in south Florida inferred from otolith shape analysis. *Mar. Coast. Fish.* 195-204.
- Shepard, K.E., W.F. Patterson, D.A. DeVries, and M. Ortiz. 2010. Contemporary versus historical estimates of king mackerel (*Scomberomorus cavalla*) age and growth in the U.S. Atlantic Ocean and Gulf of Mexico. *Bull. Mar. Sci.* 86:515-532.
- Shepherd, G.R., J. Moser, D. Deuel, and P. Carlsen. 2006. The migration patterns of bluefish (*Pomatomus saltatrix*) along the Atlantic coast determined from tag recoveries. *Fish. Bull.* 104:559-570.

- Sierra, L.M., R. Claro and O.A. Popova. 1994. Alimentacion y relaciones tróficas. p. 263–284. *In*: Rodolfo Claro (ed.) *Ecología de los Peces Marinos de Cuba*. Instituto de Oceanología Academia de Ciencias de Cuba and Centro de Investigaciones de Quintana Roo, Mexico.
- Simmons, C.M., S.T. Szedlmayer. 2011. Recruitment of age-0 gray triggerfish to benthic structured habitat in the northern Gulf of Mexico. *Trans. Amer. Fish. Soc.* 140:14-20.
- Simmons, E.G. 1957. An ecological survey of the upper Laguna Madre of Texas. *Publ. Inst. Mar. Sci.* 4:156–200.
- Sims, H.W., Jr., and R.M. Ingle. 1967. Caribbean recruitment of Florida's spiny lobster populations. *Q. J. Fla. Acad. Sci.* 29:207–242.
- Sindermann, C. J. 1979. Pollution-associated diseases and abnormalities of fish and shellfish: a review. *Fish. Bull.* 76:717-749.
- Smith, D. G. 1989. Order Elopiformes; Families Elopidae, Megalopidae, and Albulidae Leptocephali, p. 961–972. *Memoir Sears Foundation for Marine Research*. Number 1. *Fishes of the Western North Atlantic*. Part Nine. Volume Two Leptocephali. Sears Fnd. for Marine Research, Yale University.
- Smith, J.W. 1995. Life history of cobia, *Rachycentron canadum* (Osteichthyes: Rachycentridae), in North Carolina waters. *Brimleyana* 23:1–23.
- Smith, J.W., and C.A. Wenner. 1985. Biology of the southern kingfish in the South Atlantic Bight. *Trans. Amer. Fish. Soc.* 114:356–366.
- Snelson, F.F., Jr. 1992. Biological studies of the crevalle jack (*Caranx hippos*) in Florida. Final Report submitted to the Florida Dept. Environ. Protection, Florida Marine Research Institute, St. Petersburg.
- Sogard, S.M., G.V.N. Powell, and J.G. Holmquist. 1989. Utilization by fishes of shallow, seagrass-covered banks in Florida Bay: 2. Diel and tidal patterns. *Environ Biol. Fishes* 24:81–92.
- South Atlantic Fishery Management Council. 1993. Fishery management plan for the shrimp fishery of the South Atlantic region including a final environmental impact study and regulatory impact review. South Atlantic Fishery Management Council, Charleston, South Carolina.
- South Atlantic Fishery Management Council. 1998. Proceedings of the dolphin/wahoo workshop. Charleston, South Carolina.
- South Atlantic Fishery Management Council. 2010. Amendment 17A to the fishery management plan for the snapper grouper fishery of the South Atlantic region with final environmental impact statement, initial regulatory flexibility act analysis, regulatory impact review, and social impact assessment/fishery impact statement. SAFMC-NMFS, North Charleston, South Carolina. 375 p.

- South Atlantic Fishery Management Council and Gulf of Mexico Fishery Management Council. 1981. Profile of the calico scallop fishery in the South Atlantic and Gulf of Mexico. So. Atl. Fish. Manage. Council, Charleston, South Carolina.
- Springer, V.G., and K.D. Woodburn. 1960. An ecological study of the fishes of the Tampa Bay area. Fl. St. Bd. Conserv., Prof. Pap. Ser. 1:1–104.
- Stark, W.A., II. 1971. Biology of the gray snapper, *Lutjanus griseus* (Linnaeus) in the Florida Keys. Pages 11–150 in W.A. Stark and R.E. Schroeder (eds.) Investigations on the gray snapper, *Lutjanus griseus*. Trop. Oceanogr. (Miami) 10.
- Steele, P. Unpubl ms. Stock assessment profile for the penaeid shrimp fisheries of the southeastern United States and Gulf of Mexico. Florida Marine Research Institute, St. Petersburg.
- Steele, P. 1979. A synopsis of the biology of the blue crab *Callinectes sapidus* Rathbun in Florida. Proc. Blue Crab Colloquium, Oct. 18–19.
- Steele, P. 1991. Population dynamics and migration of the blue crab, *Callinectes sapidus* (Rathbun), in the eastern Gulf of Mexico. Proc. Gulf Caribb. Fish. Inst. 40:241–244.
- Steele, P., and H.M. Perry (editors). 1990. The blue crab fishery of the Gulf of Mexico United States: a regional management plan. Gulf States Marine Fisheries Commission Report 21, Ocean Springs, Ms.
- Stevens, P. W., D. A. Blewett, R. E. Boucek, J. S. Rehage, B. L. Winner, J. M. Young, J. A. Whittington, and R. Paperno. 2016. Resilience of a tropical sport fish population to a severe cold event varies across five estuaries in southern Florida. Ecosphere 7(8):e01400. 10.1002/ecs2.1400.
- Stevens, J.D. and P.D. Wiley. 1986. Biology of two commercially important carcharhinid sharks from northern Australia. Aust. J. Mar. Freshwat. Res. 37(6):671–688.
- Stickney, R.R., G.L. Taylor, and D.B. White. 1975. Food habits of five species of young southeastern United States estuarine Sciaenidae. Chesapeake Sci. 16:104–114.
- Stokes, G.M. 1977. Life history studies of southern flounder (*Paralichthys lethostigma*) and gulf flounder (*P. albigutta*) in the Aransas Bay area of Texas. Tex. Pks. Wildl. Dept. Tech Ser. 25.
- Stoner, A.W. 1979. Species-specific predation on amphipod crustacea by the pinfish, *Lagodon rhomboides*: mediation by macrophyte standing crop. Mar. Biol. 55:201–207.
- Stoner, A.W. 1980. The feeding ecology of *Lagodon rhomboides*: variation and functional response. Fish. Bull., U.S. 78:337–352.
- Stoner, A.W. 1982. The influence of benthic macrophytes on the foraging behavior of pinfish, *Lagodon rhomboides* (Linnaeus). J. Exp. Mar. Biol. Ecol. 58:271–284.

- Stunz, G.W., T.L Linton, and R.L. Colura. 2000. Age and growth of southern flounder in Texas waters, with emphasis on Matagorda Bay. *Trans. Amer. Fish. Soc.* 129:119–125.
- Sutherland, D.F., and W.A. Fable, Jr. 1980. Results of a king mackerel (*Scomberomorus cavalla*) and Atlantic Spanish mackerel (*Scomberomorus maculatus*) migration study, 1975–79. U.S. Dept. Commerce NOAA NMFS-SEFC-12. 18p.
- Sutter, F.C., B.M. Mahmoudi, R.P. Trapani. 1992. Investigations of inshore and offshore population dynamics of Spanish sardines along the central west coast of Florida. MARFIN Final Report for NOAA Grant NA90AA-H-MF086.
- Szedlmeyer, S.T. and J.C. Howe 1997. Substrate preferences in age-0 red snapper, *Lutjanus campechanus*. *Environ. Biol. Fish.* 50:203–207.
- Tabb, D.C., and R.B. Manning. 1961. A checklist of the flora and fauna of northern Florida Bay and the adjacent brackish waters of the Florida mainland collected during the period July, 1957 through September, 1960. *Bull. Mar. Sci.* 11:552–649.
- Tagatz, M.E. 1968. Biology of the blue crab, *Callinectes sapidus*, in the St. Johns River, Florida. U.S. Fish Wildl. Serv. Fish. Bull. 67:17–33.
- Tang, Q. 1983. Assessment of the blue crab commercial fishery in Chesapeake Bay. Pages 24–34, In P. Jones, D. Heimbuch, and C. Stagg (eds.), Report of the Workshop on Blue Crab Stock Dynamics in Chesapeake Bay. Chesapeake Biological Laboratory, Center for Environmental and Estuarine Studies, University of Maryland.
- Taylor, R.G., H.J. Grier , and J.A. Whittington. 1998. Spawning rhythms of common snook in Florida. *J. Fish. Biol.* 53:502–520.
- Taylor, R.G., J.A. Whittington, H.J. Grier, and R.E. Crabtree. 2000. Age, growth, maturation, and protandric sex reversal in common snook, *Centropomus undecimalis*, from the east and west coasts of Florida. *Fish. Bull.*, U.S. 98:612–624.
- Thomas, D.L. 1971. The early life history and ecology of six species of drum (Sciaenidae) in the lower Delaware River, a brackish tidal estuary. *Ichthyological Associates, Delaware Progress Report* 3(3): 247p.
- Thompson, B.A., M. Beasley, and C.A. Wilson. 1999. Age distribution and growth of greater amberjack, *Seriola dumerili*, from the north-central Gulf of Mexico. *Fish. Bull.* 97:362–371.
- Thompson, N. 1998. Characterization of the dolphin fish (Coryphaenidae, Pisces) fishery of the United States western North Atlantic ocean. Report to the Gulf of Mexico Fishery Management Mackerels Stock Assessment Panel MSAP/03 from the NOAA-NMFS Southeast Fisheries Science Center, Miami.
- Thompson, R. and J.L. Munro. 1974. The biology, ecology, and bionomics of the snappers, Lutjanidae. In: *Caribbean Coral Reef Fishery Resources*, ed. By J.L. Munro. ICLARM Studies and Reviews 7: 94-109.

- Thomson, J.M. 1963. Synopsis of biological data on the grey mullet *Mugil cephalus*, Linnaeus 1758. Australia CSIRO Division of Fisheries Oceanography Fishery Synopsis 1:68p.
- Thorrold, S.R., C.M. Jones, P.K. Swart, and T.E. Targett. 1998. Accurate classification of juvenile weakfish *Cynoscion regalis* to estuarine nursery areas based on chemical signatures of otoliths. Mar. Ecol. Prog. Ser. 173:253–265.
- Thorrold, S.R., C.L. Latkoczy, P.K. Swart, and C.M. Jones. 2001. Natal homing in a marine fish metapopulation. Science 291:297-299.
- Tinker, S.W. 1978. Fishes of Hawaii, a handbook of the marine fishes of Hawaii and the Central Pacific Ocean. Hawaiian Service Inc., Honolulu. 568p.
- Topp, R.W., and F.H. Hoff, Jr. 1972. Flatfishes (Pleuronectiformes). Mem. Hourglass Cruises Vol. 4; Part 2, 135p.
- Tringali, M.D., and T.M. Bert. 1996. The genetic stock structure of common snook, *Centropomus undecimalis*. Can. J. Fish Aquat. Sci. 53:974–984.
- Tringali, M.D., and R.R. Wilson, Jr. 1993. Differences in haplotype frequencies of mtDNA of the Spanish sardine *Sardinella aurita* between specimens from the eastern Gulf of Mexico and southern Brazil. Fish. Bull., U.S. 91:362–370.
- Turner, S.C., N.J. Cummings, and C.E. Porch. 2000. Stock assessments of Gulf of Mexico greater amberjack using data through 1998. NMFS-SEFSC Miami Doc. SFD 99/00-100.
- Tweeddale W.A., T.M. Bert, and S.D. Brown. 1993. Growth of postsettlement juveniles of the Florida stone crab, *Menippe mercenaria* (Say) (Decopod: Xanthidae), in the laboratory. Bull. Mar. Sci. 52:873–885.
- Uchiyama, J.H., R.K. Burch, and S.A. Kraul, Jr. 1986. Growth of dolphins, *Coryphaena hippurus* and *C. eqiselis*, in Hawaiian waters as determined by daily increments on otoliths. Fish. Bull., U.S. 84:186–191
- Van Engel, W.A. 1958. The blue crab and its fishery in Chesapeake Bay. Part 1, Reproduction, early development, growth and migration. Comm. Fish. Rev. 20:6–17.
- van Voorhees, D.A., T.R. Sminkey, J.W. Schlechte, D.M. Donaldson, K.J. Anson, J.R. O’Hop, M.D.B. Norris, J.A. Shepherd, T. van Devender, R.F. Zales, Jr. 2002. The new Marine Recreational Fishery Statistics Survey method for estimating charter boat fishing effort. Proc. Gulf Carib. Fisheries Inst. 53: 332-343.
- Vaughan, D.S. 1996. Status of the red drum stocks of the Atlantic coast: stock assessment report for 1995. NMFS, SEFSC Beaufort Lab. March 1996.

- Vaughan, D.S. and J.T. Carmichael. 2000. Assessment of Atlantic red drum for 1999: northern and southern regions. NOAA Technical Memorandum NMFS-SEFSC-447.
- Vaughan, D.S., and J. W. Smith 1988. A stock assessment of the Atlantic menhaden, *Brevoortia patronus*, fishery. NOAA Tech. Rep. NMFS 63, 18p.
- Vaughan, D. S., J. W. Smith, and M. H. Prager 2000. Population Characteristics of Gulf menhaden, *Brevoortia patronus*. NOAA Tech. Rep. NMFS. 149.19p.
- Vaughan, D. S., J. W. Smith, and E. H. William 2002. Analyses on the status of the Atlantic menhaden stock. Report to the Menhaden Technical Committee. 48p.
- Vaughan, D. S. 1998. Atlantic menhaden stock assessment. A report to the Atlantic State Marine Fisheries Commission Peer Review. October 29, 1998.149p.
- Wade, R.A. 1962. The biology of the tarpon, *Megalops atlanticus*, and the oxeye, *Megalops cyprinoides*, with emphasis on larval development. Bull. Mar. Sci. Gulf Caribb. 12:545–622.
- Wenner, C.A., and W.A. Roumillat, J.E. Moran, Jr., M.B. Maddox, L.B. Daniel III, and J.W. Smith. 1990. Investigations on the life history and population dynamics of marine recreational fishes in South Carolina: Part 1. SCWMRD Mar. Resour. Res. Inst.
- West, J., G. Decossas, A. Melancon, S. Potts, J. Powers 2014 *Update Assessment of Spotted Seatrout *Cynoscion nebulosus* in Louisiana Waters*. [www.wlf.louisiana.gov/sites/default/files/pdf/page/37756-stock-assessments/lasstassessment2014.pdf](http://www.wlf.louisiana.gov/sites/default/files/pdf/page/37756-stock-assessments/lasstassessment2014.pdf). Accessed 19 Sept. 2017.
- White, M.L. and M.E. Chittenden, Jr. 1977. Age determination, reproduction, and population dynamics of Atlantic croaker, *Micropogonias undulatus*. Fish. Bull., U.S. 75:109–123.
- Williams, E.H. 2001. Assessment of cobia, *Rachycentron canadum*, in the waters of the Gulf of Mexico. National Oceanic and Atmospheric Administration. Center for Coastal Fisheries and Habitat Research. Beaufort, NC
- Wilson, C.A., and D.L. Nieland. 1997. Age and size distribution of commercially harvested red snapper (*Lutjanus campechanus*) in the northern Gulf of Mexico. Tenth Annual MARFIN Conference, December 10–11, Tampa, Florida.
- Wilson, C.A., J.H. Render, and D.L. Nieland. 1994. Life history gaps in red snapper (*Lutjanus campechanus*), swordfish (*Xiphias gladius*), and red drum (*Sciaenops ocellatus*) in the northern Gulf of Mexico; age distribution, growth, and some reproductive biology. Final Report to the U.S. Dept. Comm. Nat. Mar. Fish. Serv., Marine Fisheries Initiative (MARFIN) Coop. Agreement NA17FF0383-02. 79p.
- Wilson, M.M, T. M. Bert, and S. Seyoum. 2002. Genetic stock structure of the spotted seatrout, *Cynoscion nebulosus*, in Florida. Florida Marine Research Institute Report Number IHR2002-005. 11p.
- Wood, A.D. 2013. Bluefish 2013 Stock Assessment Update. Northeast Fisheries Science Center,

National Marine Fisheries Service, Woods Hole, MA. 38 p.

Zale, A.V., and S.G. Merrifield. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (South Florida)—ladyfish and tarpon. U.S. Fish and Wildl. Serv. Biol. Rep. 82(11.104). U.S. Army Corps of Engineers, TR EL-82-4. 17p.

Zhao, B., and J.C. McGovern. 1997. Temporal variation in sexual maturity and gear-specific sex ratio of the vermilion snapper, *Rhomboplites aurorubens*, in the South Atlantic Bight. Fish. Bull. 95:838848.

Zhao, B., J.C. McGovern, and P.J. Harris. 1997. Age, growth, and temporal change in size at age of the vermilion snapper from the South Atlantic Bight. Trans. Amer. Fish. Soc. 126:181–193.